

Title	GENERALIZED HEEGAARD SPLITTINGS AND THE DISK COMPLEX
Author(s)	Kim, Jungsoo
Citation	Osaka Journal of Mathematics. 57(1) p.103-p.140
Issue Date	2020-01
oaire:version	VoR
URL	https://doi.org/10.18910/73740
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

GENERALIZED HEEGAARD SPLITTINGS AND THE DISK COMPLEX

JUNGSOO KIM

(Received January 23, 2018, revised September 4, 2018)

Abstract

Let M be an orientable, irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ a weakly reducible, unstabilized Heegaard splitting of M of genus at least three. In this article, we define an equivalence relation \sim on the set of the generalized Heegaard splittings obtained by weak reductions and find special subsets of the disk complex $\mathcal{D}(F)$ named by the “*equivalent clusters*”, where we can find a canonical function Φ from the set of equivalent clusters to the set of the equivalent classes for the relation \sim . These equivalent classes are more detailed than the isotopy classes of the generalized Heegaard splittings obtained by weak reductions from F . In the last section, we prove Φ is a bijection if the genus of F is three.

1. Introduction and Result

Let M be an orientable, irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ an unstabilized Heegaard splitting of M of genus at least three.

In this article, we will define an equivalence relation \sim on the set of the generalized Heegaard splittings obtained by weak reductions¹, where two generalized Heegaard splittings \mathbf{H}_1 and \mathbf{H}_2 obtained by weak reductions are *equivalent* or $\mathbf{H}_1 \sim \mathbf{H}_2$ if the following hold:

- (1) \mathbf{H}_1 is isotopic to \mathbf{H}_2 in M as two sets of surfaces,
- (2) $\text{Thick}(\mathbf{H}_1) \cap \mathcal{V}$ is isotopic to $\text{Thick}(\mathbf{H}_2) \cap \mathcal{V}$ in \mathcal{V} ,
- (3) $\text{Thick}(\mathbf{H}_1) \cap \mathcal{W}$ is isotopic to $\text{Thick}(\mathbf{H}_2) \cap \mathcal{W}$ in \mathcal{W} , and
- (4) $\text{Thin}(\mathbf{H}_1) \cap \text{int}(M)$ is isotopic to $\text{Thin}(\mathbf{H}_2) \cap \text{int}(M)$ in M ,

where $\text{Thick}(\mathbf{H}_i)$ and $\text{Thin}(\mathbf{H}_i)$ are the thick and thin levels of \mathbf{H}_i for $i = 1, 2$, respectively. Then we will prove the following theorem.

Theorem 1.1. *Suppose M is an orientable, irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ is a weakly reducible, unstabilized Heegaard splitting of M of genus at least three. Then there are special subsets of $\mathcal{D}(F)$ named by the “*equivalent clusters*” such that there is a canonical function Φ from the set of the equivalent clusters to the set of the equivalent classes for the relation \sim . The equivalent clusters satisfy the following:*

- (1) *Each equivalent cluster is contractible.*
- (2) *For a given equivalent cluster, the generalized Heegaard splittings obtained by weak reductions along the weak reducing pairs in the cluster are all equivalent. Moreover,*

2010 Mathematics Subject Classification. 57M50.

¹We will consider a *weak reduction* as in [1] consisting of two steps, *pre-weak reduction* and *cleaning* to discard the pairs of thick and thin levels cobounding products in the isotopy classes of the generalized Heegaard splittings.

the equivalent clusters are classified into ten types (Definition 3.19, Definition 3.22, Definition 3.23 and Definition 3.24).

- (3) If there is a simplex of $\mathcal{D}(F)$ such that the generalized Heegaard splittings obtained by weak reductions along the weak reducing pairs in the simplex are all equivalent, then this simplex belongs to a uniquely determined equivalent cluster (Theorem 3.34).

Note that the definition of the disk complex $\mathcal{D}(F)$ can be found in Section 2. In the last section, we will consider the case where the genus of F is three and prove that each component of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ is just an equivalent cluster and the canonical function Φ is bijective, where $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ is the union of all simplices of $\mathcal{D}(F)$ intersecting both $\mathcal{D}_{\mathcal{V}}(F)$ and $\mathcal{D}_{\mathcal{W}}(F)$. Using it, we will prove there is a naturally induced function Ω from the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ to the set of the isotopy classes of the generalized Heegaard splittings obtained by weak reductions and describe what Ω is.

Theorem 1.2 (Theorem 4.4). *Let $(\mathcal{V}, \mathcal{W}; F)$ be a weakly reducible, unstabilized Heegaard splitting of genus three in an orientable, irreducible 3-manifold M . Then the domain of Φ is the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$, Φ is bijective, and there is a canonically induced function Ω from the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ to the set of the isotopy classes of the generalized Heegaard splittings obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$. The number of components of the preimage of an isotopy class of Ω is the number of ways to embed the thick level contained in \mathcal{V} into \mathcal{V} (or in \mathcal{W} into \mathcal{W}). This means if we consider a generalized Heegaard splitting \mathbf{H} obtained by weak reduction from $(\mathcal{V}, \mathcal{W}; F)$, then the way to embed the thick level of \mathbf{H} contained in \mathcal{V} into \mathcal{V} determines the way to embed the thick level of \mathbf{H} contained in \mathcal{W} into \mathcal{W} up to isotopy and vice versa.*

This article is constructed as follows. First, we will classify the generalized Heegaard splittings obtained by weak reductions into five types in Lemma 3.1 simply by the shapes of the two compression bodies intersecting the inner thin level, where the *inner thin level* is the union of the components of the thin level contained in $\text{int}(M)$. Then we will find a necessary and sufficient condition that two weak reducing pairs forming a 2-simplex in $\mathcal{D}(F)$ induce equivalent generalized Heegaard splittings after weak reductions in Lemma 3.5. Using this lemma, we will reclassify the generalized Heegaard splittings obtained by weak reductions into ten types in Definition 3.14. After this reclassification, we will define the “*equivalent clusters*” and the canonical function Φ . Of course, a component of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ might not be an equivalent cluster in general (see Theorem 3.34). In the last section, we will find a necessary and sufficient condition that each component of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ is an equivalent cluster in Lemma 4.2. Next, we will find a sufficient condition that two equivalent clusters corresponding to the same equivalent class by Φ are the same in Lemma 4.3. By applying Lemma 4.3 to the genus three case, we will prove Theorem 4.4.

2. Preliminaries

A *compression body* is a 3-manifold which can be obtained by starting with a closed, orientable, connected surface F , forming the product $F \times I$, attaching some number of 2-handles to $F \times \{1\}$ and capping off all resulting 2-sphere boundary components that are not contained in $F \times \{0\}$ with 3-balls. The boundary component $F \times \{0\}$ is referred to as ∂_+ and the

rest of the boundary is referred to as ∂_- . For a compression body \mathcal{V} with $\partial_- \mathcal{V} = \emptyset$, we call it a *handlebody*. For a compression body \mathcal{V} , we call the genus of $\partial_+ \mathcal{V}$ the *genus* of \mathcal{V} . For a compression body \mathcal{V} with $\partial_- \mathcal{V} \neq \emptyset$, we can obtain \mathcal{V} from $\partial_- \mathcal{V} \times I$ by attaching 1-handles to $\partial_- \mathcal{V} \times \{1\}$. This means the genus of $\partial_+ \mathcal{V}$ is at least the sum of genera of the components of $\partial_- \mathcal{V}$. For a surface F , if there is an embedded disk D in M such that $\partial D \subset F$ is an essential curve in F and $\text{int}(D) \cap F = \emptyset$, then we call D a *compressing disk* for F . For a compression body \mathcal{V} , if D is properly embedded in \mathcal{V} and it is a compressing disk for $\partial_+ \mathcal{V}$, then we call D a *compressing disk* in \mathcal{V} . A *Heegaard splitting* of a 3-manifold M is an expression of M as a union $\mathcal{V} \cup_F \mathcal{W}$, denoted as $(\mathcal{V}, \mathcal{W}; F)$, where \mathcal{V} and \mathcal{W} are compression bodies that intersect in a transversally oriented surface $F = \partial_+ \mathcal{V} = \partial_+ \mathcal{W}$. We say F is the *Heegaard surface* of this splitting. If \mathcal{V} or \mathcal{W} is homeomorphic to a product, then we call such compression body *trivial* and we say the splitting is *trivial*. If there are compressing disks $V \subset \mathcal{V}$ and $W \subset \mathcal{W}$ such that $V \cap W = \emptyset$, then we say the splitting is *weakly reducible* and call the pair (V, W) a *weak reducing pair*. If (V, W) is a weak reducing pair and ∂V is isotopic to ∂W in F , then we call (V, W) a *reducing pair*. If there is a pair of compressing disks (\bar{V}, \bar{W}) such that \bar{V} intersects \bar{W} transversely in a point in F , then we call this pair a *canceling pair* and say the splitting is *stabilized*. Otherwise, we say the splitting is *unstabilized*.

Let F be a surface of genus at least two in a compact, orientable 3-manifold M . Then the *disk complex* $\mathcal{D}(F)$ is defined as follows:

- (i) Vertices of $\mathcal{D}(F)$ are isotopy classes of compressing disks for F .
- (ii) A set of $m + 1$ vertices forms an m -simplex if there are representatives for each that are pairwise disjoint.

Consider a Heegaard splitting $(\mathcal{V}, \mathcal{W}; F)$ of genus at least two in an orientable, irreducible 3-manifold M . Let $\mathcal{D}_{\mathcal{V}}(F)$ and $\mathcal{D}_{\mathcal{W}}(F)$ be the subspaces of $\mathcal{D}(F)$ spanned by compressing disks in \mathcal{V} and \mathcal{W} , respectively. We call these subspaces *the disk complexes of \mathcal{V} and \mathcal{W}* , respectively. Let $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ be the subset of $\mathcal{D}(F)$ consisting of all simplices with at least one vertex from $\mathcal{D}_{\mathcal{V}}(F)$ and at least one vertex from $\mathcal{D}_{\mathcal{W}}(F)$. By definition, if there is a simplex σ in $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ such that $\sigma \subset \mathcal{D}_{\mathcal{V}}(F)$ or $\mathcal{D}_{\mathcal{W}}(F)$, then it must be a proper subsimplex of a simplex intersecting both $\mathcal{D}_{\mathcal{V}}(F)$ and $\mathcal{D}_{\mathcal{W}}(F)$.

If there is no confusion, we will use the disk V instead of the isotopy class $[V] \in \mathcal{D}(F)$ for the sake of convenience. Hence, we will say (1) $D_1 = D_2$ in $\mathcal{D}(F)$ instead of $[D_1] = [D_2]$ and (2) if there is a k -simplex Δ in $\mathcal{D}(F)$, then we will denote Δ as a set of $k + 1$ mutually disjoint, non-isotopic compressing disks for F . Note that if there is a compressing disk in a compression body \mathcal{V} , then the isotopy class of the boundary of the disk in $\partial_+ \mathcal{V}$ determines the isotopy class of the compressing disk in \mathcal{V} . (Since $\partial F = \emptyset$, the pair (\mathcal{V}, F) is *spotless*. Therefore, the natural map $\nu : \mathcal{D}_{\mathcal{V}}(F) \rightarrow \mathcal{C}(F)$ induced by taking a compressing disk in \mathcal{V} to its boundary is injective, where $\mathcal{C}(F)$ is the *curve complex* of F , see [7] for the details.) This gives an important lemma.

Lemma 2.1. *Let \mathcal{V} be a compression body. If there are two compressing disks V_1 and V_2 in \mathcal{V} such that ∂V_1 is isotopic to ∂V_2 in $\partial_+ \mathcal{V}$, then V_1 is isotopic to V_2 in \mathcal{V} .*

We get the following corollary directly from Lemma 2.1.

Corollary 2.2. *Let \mathcal{V} be a compression body and $\mathcal{V}' \subset \mathcal{V}$ a compression body such that $\partial_+ \mathcal{V} = \partial_+ \mathcal{V}'$. If V_1 and V_2 are compressing disks in \mathcal{V}' such that V_1 is isotopic to V_2 in \mathcal{V}' ,*

then V_1 is isotopic to V_2 in \mathcal{V} .

Note that $D_{\mathcal{V}}(F)$ and $D_{\mathcal{W}}(F)$ are contractible (see [8]). Let $(\mathcal{V}, \mathcal{W}; F)$ be a Heegaard splitting of genus at least two. Since $\dim(C(F))$ is finite and there exist at most two disks in $D(F)$ which correspond to an element of $C(F)$ (such two disks exist if and only if there exists a reducing pair), $\dim(D(F))$ is also finite.

From now on, we will consider only unstabilized Heegaard splittings of an irreducible 3-manifold. If a Heegaard splitting of a compact 3-manifold is reducible, then the manifold is reducible or the splitting is stabilized (see [11]). Hence, we can exclude the possibilities of reducing pairs among weak reducing pairs.

Suppose W is a compressing disk for a properly embedded surface $F \subset M$. Then there is a subset of M that can be identified with $W \times I$ so that $W = W \times \{\frac{1}{2}\}$ and $F \cap (W \times I) = (\partial W) \times I$. We form the surface F_W , obtained by *compressing F along W* , by removing $(\partial W) \times I$ from F and replacing it with $W \times (\partial I)$. We say the two disks $W \times (\partial I)$ in F_W are the *scars* of W .

Lemma 2.3 (Lustig and Moriah, Lemma 1.1 of [6]). *Suppose M is an irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ is an unstabilized Heegaard splitting of M . If F' is obtained by compressing F along a collection of pairwise disjoint disks, then no S^2 component of F' can have scars from disks in both \mathcal{V} and \mathcal{W} .*

The next is the definition of “generalized Heegaard splitting” originated from [13].

DEFINITION 2.4 (DEFINITION 4.1 OF [1]). A *generalized Heegaard splitting* (we will use *GHS* as abbreviation from now on) \mathbf{H} of a 3-manifold M is a pair of sets of pairwise disjoint, transversally oriented, connected surfaces, $\text{Thick}(\mathbf{H})$ and $\text{Thin}(\mathbf{H})$ (resp. called the *thick levels* and *thin levels*), which satisfies the following conditions.

- (1) Each component M' of $M - \text{Thin}(\mathbf{H})$ meets a unique element H_+ of $\text{Thick}(\mathbf{H})$ and H_+ is a Heegaard surface in the closure of M' , say $M(H_+)$.
- (2) As each Heegaard surface $H_+ \subset M(H_+)$ is transversally oriented, we can consistently talk about the points of $M(H_+)$ that are “above” H_+ or “below” H_+ . Suppose $H_- \in \text{Thin}(\mathbf{H})$. Let $M(H_+)$ and $M(H'_+)$ be the submanifolds on each side of H_- . Then H_- is below H_+ if and only if it is above H'_+ .
- (3) There is a partial ordering on the elements of $\text{Thin}(\mathbf{H})$ which satisfies the following: Suppose H_+ is an element of $\text{Thick}(\mathbf{H})$, H_- is a component of $\partial M(H_+)$ above H_+ and H'_- is a component of $\partial M(H_+)$ below H_+ . Then $H_- > H'_-$.

We denote the maximal subset of $\text{Thin}(\mathbf{H})$ consisting of surfaces in $\text{int}(M)$ as $\overline{\text{Thin}}(\mathbf{H})$ and call it the *inner thin level*.

DEFINITION 2.5 (A RESTRICTED VERSION OF DEFINITION 5.2, DEFINITION 5.3, AND DEFINITION 5.6 OF [1]). Let M be an orientable, irreducible 3-manifold. Let \mathbf{H} be an unstabilized Heegaard splitting of M , i.e. $\text{Thick}(\mathbf{H}) = \{F\}$ and $\text{Thin}(\mathbf{H})$ consists of ∂M . Let V and W be disjoint compressing disks for F from the opposite sides of F . Here, Lemma 2.3 guarantees F_{VW} does not have a 2-sphere component. Define

$$\text{Thick}(\mathbf{G}') = (\text{Thick}(\mathbf{H}) - \{F\}) \cup \{F_V, F_W\}, \text{ and}$$

$$\text{Thin}(\mathbf{G}') = \text{Thin}(\mathbf{H}) \cup \{F_{VW}\},$$

where we assume each element of $\text{Thick}(\mathbf{G}')$ belongs to $\text{int}(\mathcal{V})$ or $\text{int}(\mathcal{W})$ by slightly pushing F_V and F_W off into $\text{int}(\mathcal{V})$ and $\text{int}(\mathcal{W})$, respectively, and then also assume they miss F_{VW} . (In the proof of Lemma 2.16 of [5], the author used the term F_V for denoting $\text{Thick}(\mathbf{G}') \cap \mathcal{V}$ for the sake of convenience. In this article, we will only use this term for this definition and we will use \tilde{F}_V for general cases to avoid confusion.) We say the GHS $\mathbf{G}' = \{\text{Thick}(\mathbf{G}'), \text{Thin}(\mathbf{G}')\}$ is obtained from \mathbf{H} by *preweak reduction* along (V, W) . We can see F_V comes first, F_{VW} next, and F_W last in M or vice versa (see Figure 10 of [1]). If there are elements $S \in \text{Thick}(\mathbf{G}')$ and $s \in \overline{\text{Thin}}(\mathbf{G}')$ that cobound a product region P of M such that $P \cap \text{Thick}(\mathbf{G}') = S$ and $P \cap \text{Thin}(\mathbf{G}') = s$, then remove S from $\text{Thick}(\mathbf{G}')$ and s from $\text{Thin}(\mathbf{G}')$. If we repeat this procedure until there are no such two elements of $\text{Thick}(\mathbf{G}')$ and $\overline{\text{Thin}}(\mathbf{G}')$, then we get the resulting GHS \mathbf{G} of M from the GHS \mathbf{G}' (see Lemma 5.4 of [1]) and we say \mathbf{G} is obtained from \mathbf{G}' by *cleaning*. We say the GHS \mathbf{G} of M given by preweak reduction along (V, W) , followed by cleaning, is obtained from \mathbf{H} by *weak reduction* along (V, W) . From now on, we will denote the GHS \mathbf{G} obtained by weak reduction from the Heegaard surface F along a weak reducing pair (V, W) as the ordered triple $(\bar{F}_V, \bar{F}_{VW}, \bar{F}_W)$ without denoting ∂M for the sake of convenience, where (i) $\text{Thick}(\mathbf{G}) \cap \mathcal{V} = \bar{F}_V$, $\text{Thick}(\mathbf{G}) \cap \mathcal{W} = \bar{F}_W$, and $\overline{\text{Thin}}(\mathbf{G}) = \bar{F}_{VW}$ and (ii) the notations \bar{F}_V , \bar{F}_W and \bar{F}_{VW} imply they come from F_V , F_W and F_{VW} , respectively. Note that we will confirm the GHSs obtained by weak reductions in Lemma 3.1.

In this article, we will only consider “*single weak reductions*” from F , i.e. we do not perform twice or more weak reductions. Hence, “*GHSs obtained by weak reductions from F* ” means that each GHS of them consists of exactly four compression bodies (see Lemma 3.1).

DEFINITION 2.6. Let F be a weakly reducible Heegaard surface in an orientable, irreducible 3-manifold M . We say the GHSs \mathbf{H}_1 and \mathbf{H}_2 are *isotopic* if there is an ambient isotopy h_t defined on M such that h_t sends $\text{Thick}(\mathbf{H}_1) \cup \text{Thin}(\mathbf{H}_1)$ to $\text{Thick}(\mathbf{H}_2) \cup \text{Thin}(\mathbf{H}_2)$. Let \mathcal{GHS}_F be the set of isotopy classes of the GHSs obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$. We will say two GHSs $\mathbf{H}_1 = (\bar{F}_{V_1}, \bar{F}_{V_1W_1}, \bar{F}_{W_1})$ and $\mathbf{H}_2 = (\bar{F}_{V_2}, \bar{F}_{V_2W_2}, \bar{F}_{W_2})$ obtained by weak reductions are *equivalent* if (i) \mathbf{H}_1 is isotopic to \mathbf{H}_2 , i.e. $[\mathbf{H}_1] = [\mathbf{H}_2]$ in \mathcal{GHS}_F (ii) \bar{F}_{V_1} is isotopic to \bar{F}_{V_2} in \mathcal{V} , (iii) \bar{F}_{W_1} is isotopic to \bar{F}_{W_2} in \mathcal{W} , and (iv) $\bar{F}_{V_1W_1}$ is isotopic to $\bar{F}_{V_2W_2}$ in M .

We get the following lemma from Definition 2.6 and omit the trivial proof.

Lemma 2.7. Let \mathcal{GHS}_F^* be the set of GHSs obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$ and \sim the relation defined on \mathcal{GHS}_F^* such that $\mathbf{H}_1 \sim \mathbf{H}_2$ if \mathbf{H}_1 and \mathbf{H}_2 are equivalent. Then \sim is an equivalence relation on \mathcal{GHS}_F^* .

Considering Lemma 2.7, \mathcal{GHS}_F^*/\sim is the set of the equivalent classes. We will denote the equivalent class of a GHS $\mathbf{H} \in \mathcal{GHS}_F^*$ as $[\mathbf{H}]^*$ to distinguish it from the isotopy class $[\mathbf{H}]$.

Note that for any $[\mathbf{H}] \in \mathcal{GHS}_F$ (resp. $[\mathbf{H}]^* \in \mathcal{GHS}_F^*/\sim$), no pair of thick and thin levels cobounds a product region by “*cleaning*” in Definition 2.5. This means the thin level in the GHS after weak reduction may be connected even if that in the GHS after pre-weak reduction is disconnected (refer to Figure 3 of [5]).

The next two lemmas give natural, but important observations for a genus $n \geq 2$ compression body having a special compressing disk such that it divides the compression body into one or two products.

Lemma 2.8 (Ido, Jang and Kobayashi, Lemma 3.3 of [2]). *Let \mathcal{V} be a genus $n \geq 2$ compression body such that $\partial_- \mathcal{V}$ consists of a genus $n - 1$ connected surface. Then there is a unique nonseparating compressing disk in \mathcal{V} up to isotopy. Since any separating compressing disk in \mathcal{V} cuts off a uniquely determined solid torus from \mathcal{V} corresponding to the disk, every nonseparating compressing disk in \mathcal{V} is isotopic to a meridian disk of the solid torus in \mathcal{V} .*

Note that the phrase “any separating compressing disk in \mathcal{V} cuts off a uniquely determined solid torus from \mathcal{V} corresponding to the disk” of Lemma 2.8 is guaranteed by Lemma 1.3 of [12] by considering the genus of each component. Using the similar idea of Lemma 2.8, we can prove the following lemma (we omit the proof).

Lemma 2.9. *Let \mathcal{V} be a genus $n \geq 2$ compression body such that $\partial_- \mathcal{V}$ consists of a genus $m \geq 1$ connected surface and a genus $(n - m) \geq 1$ connected surface. Then there is a unique compressing disk in \mathcal{V} up to isotopy and it is separating in \mathcal{V} .*

Lemma 2.10. *Let \mathcal{V} be a genus $n \geq 2$ compression body, V a compressing disk in \mathcal{V} , and $F = \partial_+ \mathcal{V}$. Let \tilde{F}_V be the surface obtained by pushing F_V off slightly into $\text{int}(\mathcal{V})$. Let \tilde{F}'_V be a component of \tilde{F}_V and $\tilde{\mathcal{V}}'$ the closure of the component of $\mathcal{V} - \tilde{F}'_V$ intersecting F . Then $\tilde{\mathcal{V}}'$ is a compression body such that $\partial_+ \tilde{\mathcal{V}}' = \partial_+ \mathcal{V}$.*

Proof. If V is nonseparating, then $\tilde{\mathcal{V}}'$ is obtained from $\tilde{F}_V \times I$ by attaching a 1-handle whose cocore disk is V to $\tilde{F}_V \times \{1\}$, i.e. $\tilde{\mathcal{V}}'$ is a compression body.

Suppose V is separating in \mathcal{V} . Let \mathcal{V}' and \mathcal{V}'' be the 3-manifolds obtained by cutting \mathcal{V} along a small product neighborhood of V in \mathcal{V} and assume \mathcal{V}' contains \tilde{F}'_V . Then \mathcal{V}' and \mathcal{V}'' are compression bodies such that $\partial_- \mathcal{V}' \cup \partial_- \mathcal{V}'' = \partial_- \mathcal{V}$ by Lemma 1.3 of [12]. Moreover, the region between \tilde{F}'_V and $\partial_+ \mathcal{V}'$ in \mathcal{V}' is the trivial compression body $\tilde{F}'_V \times I$ and say $\partial_-(\tilde{F}'_V \times I) = \tilde{F}'_V$. Then $\tilde{\mathcal{V}}'$ is obtained from the union of $\tilde{F}'_V \times I$ and \mathcal{V}'' by attaching a 1-handle connecting $\partial_+(\tilde{F}'_V \times I)$ and $\partial_+ \mathcal{V}''$ whose cocore disk is V , i.e. $\tilde{\mathcal{V}}'$ is a compression body.

This completes the proof. \square

Lemma 2.11. *Let \mathcal{V} be a compression body of genus $n \geq 3$, V_1 and V_2 compressing disks in \mathcal{V} , and $F = \partial_+ \mathcal{V}$. Let \tilde{F}_{V_1} and \tilde{F}_{V_2} be the surfaces obtained by pushing F_{V_1} and F_{V_2} slightly into $\text{int}(\mathcal{V})$, respectively. Then the following hold:*

- (1) *Suppose both V_1 and V_2 are nonseparating in \mathcal{V} . Then \tilde{F}_{V_2} is isotopic to \tilde{F}_{V_1} in \mathcal{V} if and only if V_2 is isotopic to V_1 in \mathcal{V} .*
- (2) *Suppose V_1 is separating and V_2 is nonseparating in \mathcal{V} . Then \tilde{F}_{V_2} is isotopic to a component of \tilde{F}_{V_1} in \mathcal{V} if and only if V_1 cuts off a solid torus from \mathcal{V} and V_2 is isotopic to a meridian disk of the solid torus in \mathcal{V} .*

Proof.

- (1) Suppose both V_1 and V_2 are nonseparating in \mathcal{V} .

(\Leftarrow) We can see the region $\tilde{\mathcal{V}}$ cobounded by F_{V_2} and \tilde{F}_{V_2} is a product and we will




Fig. 1. The isotopy taking \tilde{F}_{V_2} into \tilde{F}_{V_1}' .

observe the images of $\tilde{\mathcal{V}}$ of isotopies defined on \mathcal{V} . Assume V_2 is isotopic to V_1 in \mathcal{V} by an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$, i.e. $h_1(V_2) = V_1$. After the isotopy h_t , we push off $h_1(\tilde{F}_{V_2})$ into a small neighborhood of $F \cup h_1(V_2)$ in \mathcal{V} preserving $h_1(V_2)$ by an isotopy g_t defined on \mathcal{V} (we can refer to (b) and (c) of Fig.1 ignoring the separating V_1 in the figure). Then we can assume $g_1(h_1(\tilde{F}_{V_2}))$ is \tilde{F}_{V_1}' , leading to the result.

(\Rightarrow) Assume \tilde{F}_{V_2} is isotopic to \tilde{F}_{V_1} in \mathcal{V} by an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$. Let $\tilde{\mathcal{V}}$ be the closure of the component of $\mathcal{V} - \tilde{F}_{V_1}$ intersecting F . Then $\tilde{\mathcal{V}}$ is a genus n compression body with negative boundary consisting of a genus $(n - 1)$ surface by Lemma 2.10, where $\partial_+ \tilde{\mathcal{V}} = \partial_+ \mathcal{V}$. Here, (i) $h_1(V_2)$ and V_1 are properly embedded in $\tilde{\mathcal{V}}$ and (ii) $\partial h_1(V_2)$ and ∂V_1 are nonseparating curves in $F = \partial_+ \tilde{\mathcal{V}}$, i.e. both $h_1(V_2)$ and V_1 are nonseparating compressing disks in $\tilde{\mathcal{V}}$. This means $h_1(V_2)$ is isotopic to V_1 in $\tilde{\mathcal{V}}$ by Lemma 2.8 and therefore $h_1(V_2)$ is isotopic to V_1 in \mathcal{V} by an isotopy g_t defined on \mathcal{V} by Corollary 2.2. Hence, the sequence of isotopies consisting of h_t and g_t realizes the isotopy from V_2 to V_1 in \mathcal{V} .

(2) Suppose V_1 is separating and V_2 is nonseparating in \mathcal{V} .

(\Leftarrow) Let $\tilde{\mathcal{V}}$ be as in the (\Leftarrow) part of the proof of (2.11) (see (a) of Fig.1). Assume V_1 cuts off a solid torus from \mathcal{V} and there is an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that $h_1(V_2)$ is a meridian disk of the solid torus. Then we can also assume $h_1(V_2)$ misses V_1 without loss of generality (see (b) of Fig.1). Using an additional isotopy $g_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that it preserves $h_1(V_2)$ and pushes $h_1(\tilde{F}_{V_2})$ into a small neighborhood of $F \cup h_1(V_2)$ in \mathcal{V} , we can assume $g_1(h_1(\tilde{\mathcal{V}}))$ is a sufficiently thin product (see (c) of Fig.1). Let \tilde{F}_{V_1}' be the genus $(n - 1)$ -component of \tilde{F}_{V_1} . Then we can see $g_1(h_1(\tilde{F}_{V_2}))$ is isotopic to \tilde{F}_{V_1}' in \mathcal{V} by an isotopy f_t defined on \mathcal{V} , where the reason is that there is a 3-ball in \mathcal{V} bounded by the S^2 consisting of a disk in $\text{int}(V_1)$ and a disk in $g_1(h_1(\tilde{F}_{V_2}))$ such that these two disks share the common boundary and therefore we can push $g_1(h_1(\tilde{F}_{V_2}))$ into the interior of the component of $\mathcal{V} - V_1$ containing \tilde{F}_{V_1}' through the 3-ball (see (c) and (d) of Fig.1). Hence, the sequence of isotopies consisting of h_t , g_t and f_t realizes the isotopy from \tilde{F}_{V_2} to \tilde{F}_{V_1}' in \mathcal{V} .

(\Rightarrow) Let $\tilde{\mathcal{V}}'$ be the region cobounded by F and \tilde{F}_{V_2} in \mathcal{V} . Then $\tilde{\mathcal{V}}'$ is a genus n

compression body with negative boundary consisting of the genus $(n - 1)$ surface \tilde{F}_{V_2} by Lemma 2.10, where $\partial_+ \tilde{\mathcal{V}}' = \partial_+ \mathcal{V}$. Assume there is an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that $h_1(\tilde{F}_{V_2}) = \tilde{F}'_{V_1}$ for a component \tilde{F}'_{V_1} of \tilde{F}_{V_1} . Then $\tilde{\mathcal{V}} = h_1(\tilde{\mathcal{V}}')$ is also a compression body homeomorphic to $\tilde{\mathcal{V}}'$, where it is the region cobounded by F and \tilde{F}'_{V_1} in \mathcal{V} . Therefore, the genus of $\partial_- \tilde{\mathcal{V}} = \tilde{F}'_{V_1}$ must be $(n - 1)$. Moreover, considering ∂V_1 and $\partial h_1(V_2)$ in $F = \partial_+ \tilde{\mathcal{V}}$, V_1 is a separating compressing disk and $h_1(V_2)$ is a nonseparating compressing disk in $\tilde{\mathcal{V}}$. This means $h_1(V_2)$ is isotopic to a meridian disk D of the solid torus that V_1 cuts off from $\tilde{\mathcal{V}}$ in $\tilde{\mathcal{V}}$ by Lemma 2.8. Therefore, $h_1(V_2)$ is isotopic to D in \mathcal{V} by an isotopy g_t defined on \mathcal{V} by Corollary 2.2. Hence, the sequence of isotopies consisting of h_t and g_t realizes the isotopy from V_2 to D in \mathcal{V} .

This completes the proof of Lemma 2.11. \square

Lemma 2.12. *Let \mathcal{V} be a compression body of genus $n \geq 2$ and V_1 and V_2 separating compressing disks in \mathcal{V} such that the geometric intersection number $i(\partial V_1, \partial V_2) = 0$ in $\partial_+ \mathcal{V}$. If each of V_1 and V_2 cuts off $S \times I$ from \mathcal{V} for a component S of $\partial_- \mathcal{V}$ such that S is the common 0-level, then V_1 is isotopic to V_2 in \mathcal{V} .*

Proof. Since the geometric intersection number $i(\partial V_1, \partial V_2) = 0$ in $\partial_+ \mathcal{V}$, moving the part of V_1 near $\partial_+ \mathcal{V}$ continuously, there is an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that $\partial h_1(V_1) \cap \partial V_2 = \emptyset$. Moreover, we can assume that $h_1(V_1)$ intersects V_2 transversely and therefore each component of $h_1(V_1) \cap V_2$ is an inessential loop in both $\text{int}(h_1(V_1))$ and $\text{int}(V_2)$. This means we can find a compressing disk V in \mathcal{V} such that $\partial V = \partial h_1(V_1)$ and $V \cap V_2 = \emptyset$ by using innermost disk arguments (see the proof of Claim 1 of Lemma 3.1.5 of [11]). Therefore, Lemma 2.1 implies $h_1(V_1)$ is isotopic to V in \mathcal{V} by an isotopy g_t defined on \mathcal{V} , i.e. $g_1(h_1(V_1)) \cap V_2 = \emptyset$. Let \mathcal{V}_1 and \mathcal{V}_2 be the $S \times I$ s that $g_1(h_1(V_1))$ and V_2 cut off from \mathcal{V} , respectively, guaranteed by the assumption of this lemma. Considering the 1-level of \mathcal{V}_1 (resp. \mathcal{V}_2), it consists of a once-punctured surface F_1 (resp. F_2) in $\partial_+ \mathcal{V}$ and $g_1(h_1(V_1))$ (resp. V_2), where $F_1 \cap g_1(h_1(V_1)) = \partial F_1 = \partial g_1(h_1(V_1))$, $F_2 \cap V_2 = \partial F_2 = \partial V_2$. Since \mathcal{V}_1 shares the 0-level S with \mathcal{V}_2 , the assumption $g_1(h_1(V_1)) \cap V_2 = \emptyset$ implies one of \mathcal{V}_1 and \mathcal{V}_2 contains the other. Assume $\mathcal{V}_1 \subset \mathcal{V}_2$ without loss of generality, i.e. $F_1 \subset F_2$. Since $g(F_1) = g(F_2) = g(S) \geq 1$, ∂F_1 cuts off a uniquely determined planar surface from F_2 . Moreover, $g_1(h_1(V_1))$ is a compressing disk in \mathcal{V} and therefore $\partial F_1 = \partial g_1(h_1(V_1))$ cannot bound a disk in F_2 . This means ∂F_1 cuts off an annulus from F_2 , i.e. $\partial F_1 (= \partial g_1(h_1(V_1)))$ is isotopic to $\partial F_2 (= \partial V_2)$ in $F_2 \subset F$. Therefore, $g_1(h_1(V_1))$ is isotopic to V_2 in \mathcal{V} by an isotopy $f_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ by Lemma 2.1. Hence, the sequence of isotopies consisting of h_t , g_t and f_t realizes the isotopy from V_1 to V_2 in \mathcal{V} .

This completes the proof. \square

Lemma 2.13. *Let \mathcal{V} be a compression body of genus $n \geq 3$, V_1 and V_2 mutually disjoint separating compressing disks in \mathcal{V} , and $F = \partial_+ \mathcal{V}$. Let \tilde{F}_{V_1} and \tilde{F}_{V_2} be the surfaces obtained by pushing F_{V_1} and F_{V_2} slightly into $\text{int}(\mathcal{V})$. Then at least one component of \tilde{F}_{V_2} is isotopic to a component of \tilde{F}_{V_1} in \mathcal{V} if and only if V_2 is isotopic to V_1 in \mathcal{V} .*

Proof. (\Leftarrow) Assume that V_2 is isotopic to V_1 in \mathcal{V} . Then considering the argument in the (\Leftarrow) part of the proof of (2.11) of Lemma 2.11, we conclude \tilde{F}_{V_2} is isotopic to \tilde{F}_{V_1} in \mathcal{V} .

(\Rightarrow) Let $\tilde{F}_{V_1} = \tilde{F}'_{V_1} \cup \tilde{F}''_{V_1}$ and $\tilde{F}_{V_2} = \tilde{F}'_{V_2} \cup \tilde{F}''_{V_2}$ by denoting each \tilde{F}_{V_i} as the union of two components for $i = 1, 2$. Assume \tilde{F}'_{V_2} is isotopic to \tilde{F}'_{V_1} by an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$. Let $\tilde{\mathcal{V}}$ be the closure of the component of $\mathcal{V} - \tilde{F}'_{V_1}$ intersecting F . Then $\tilde{\mathcal{V}}$ is a genus n compression body by Lemma 2.10, where $\partial_+ \tilde{\mathcal{V}} = \partial_+ \mathcal{V}$, and V_1 and $h_1(V_2)$ are compressing disks in $\tilde{\mathcal{V}}$. By the assumption that $V_1 \cap V_2 = \emptyset$, the geometric intersection number $i(\partial V_1, \partial h_1(V_2)) = 0$ in F . Here, we can see the product region cobounded by \tilde{F}'_{V_2} , V_2 and a subsurface $F_0 \subset F$ in \mathcal{V} has been isotoped into the region cobounded by \tilde{F}'_{V_1} , $h_1(V_2)$ and $h_1(F_0) \subset F$ in \mathcal{V} and therefore the latter region is also homeomorphic to a product. Since $\partial_+ \tilde{\mathcal{V}} = \partial_+ \mathcal{V} = F$, $\tilde{F}'_{V_1} \subset \partial_- \tilde{\mathcal{V}}$, and each of V_1 and $h_1(V_2)$ cuts off $\tilde{F}'_{V_1} \times I$ from $\tilde{\mathcal{V}}$ such that the one corresponding to V_1 shares the common 0-level \tilde{F}'_{V_1} with that corresponding to $h_1(V_2)$, Lemma 2.12 implies $h_1(V_2)$ is isotopic to V_1 in $\tilde{\mathcal{V}}$. Therefore, $h_1(V_2)$ is isotopic to V_1 by an isotopy $g_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ by Corollary 2.2. Hence, the sequence of isotopies consisting of h_t and g_t realizes the isotopy from V_2 to V_1 in \mathcal{V} .

This completes the proof. \square

Lemma 2.14. *Let F be a closed surface embedded in a 3-manifold. Let V and W be mutually disjoint compressing disks for F such that ∂V is separating and ∂W is nonseparating in F . Then ∂W is nonseparating in the relevant component of F_V .*

Proof. Let F'_V be the component of F_V containing ∂W . For the sake of contradiction, assume ∂W is separating in F'_V . Then one component of $F'_V - \partial W$ has a scar of V and the other doesn't because there is only one scar of V in F'_V by the assumption that ∂V is separating in F . This means ∂W cuts off the latter component from F , violating the assumption that ∂W is nonseparating in F . This completes the proof. \square

3. The proof of Theorem 1.1

Lemma 3.1 characterizes the possible GHSs obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$ into five types, where the five types depend on the shapes of the compression bodies intersecting the inner thin level.

Lemma 3.1 (the generalization of Appendix of [5]). *Suppose M is an irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ is a weakly reducible, unstabilized Heegaard splitting of M of genus $n \geq 3$. Let \mathbf{H} be the GHS obtained by weak reduction along a weak reducing pair (V, W) from $(\mathcal{V}, \mathcal{W}; F)$. Then \mathbf{H} is of the form $(\mathcal{V}_1, \mathcal{V}_2; \tilde{F}_V) \cup_{\tilde{F}_{VW}} (\mathcal{W}_1, \mathcal{W}_2; \tilde{F}_W)$ for four compression bodies \mathcal{V}_1 , \mathcal{V}_2 , \mathcal{W}_1 and \mathcal{W}_2 , where $\partial_- \mathcal{V}_2 \cap \partial_- \mathcal{W}_1 = \tilde{F}_{VW}$, $\tilde{F}_V \subset \mathcal{V}$, and $\tilde{F}_W \subset \mathcal{W}$. Moreover, we can classify \mathbf{H} into the following five types ((b) is divided into two types) by the shapes of $\partial_- \mathcal{V}_2$ and $\partial_- \mathcal{W}_1$ (see Fig.2).*

(a) *Each of $\partial_- \mathcal{V}_2$ and $\partial_- \mathcal{W}_1$ is connected, where either*

- (i) *V and W are nonseparating in \mathcal{V} and \mathcal{W} , respectively, and $\partial V \cup \partial W$ is nonseparating in F ,*
- (ii) *V cuts off a handlebody \mathcal{V}' of genus at least one from \mathcal{V} such that $\partial W \cap \mathcal{V}' = \emptyset$ and W is nonseparating in \mathcal{W} ,*
- (iii) *W cuts off a handlebody \mathcal{W}' of genus at least one from \mathcal{W} such that $\partial V \cap \mathcal{W}' = \emptyset$ and V is nonseparating in \mathcal{V} , or*

- (iv) V and W cut off handlebodies \mathcal{V}' and \mathcal{W}' of genus at least one from \mathcal{V} and \mathcal{W} , respectively, such that $\partial V \cap \mathcal{W}' = \emptyset$ and $\partial W \cap \mathcal{V}' = \emptyset$.

We call it a “type (a) GHS”.

- (b) One of $\partial_- \mathcal{V}_2$ and $\partial_- \mathcal{W}_1$ is connected and the other is disconnected, where either
- (i) ($\partial_- \mathcal{V}_2$ is connected) V cuts off a compression body \mathcal{V}' of genus at least one from \mathcal{V} such that $\partial_- \mathcal{V}' \neq \emptyset$ and $\partial W \cap \mathcal{V}' = \emptyset$ and W is nonseparating in \mathcal{W} ,
 - (ii) ($\partial_- \mathcal{V}_2$ is connected) V cuts off a compression body \mathcal{V}' of genus at least one from \mathcal{V} and W cuts off a handlebody \mathcal{W}' of genus at least one from \mathcal{W} such that $\partial_- \mathcal{V}' \neq \emptyset$, $\partial V \cap \mathcal{W}' = \emptyset$, and $\partial W \cap \mathcal{V}' = \emptyset$,
 - (iii) ($\partial_- \mathcal{W}_1$ is connected) W cuts off a compression body \mathcal{W}' of genus at least one from \mathcal{W} such that $\partial_- \mathcal{W}' \neq \emptyset$, $\partial V \cap \mathcal{W}' = \emptyset$, and V is nonseparating in \mathcal{V} , or
 - (iv) ($\partial_- \mathcal{W}_1$ is connected) W cuts off a compression body \mathcal{W}' of genus at least one from \mathcal{W} and V cuts off a handlebody \mathcal{V}' of genus at least one from \mathcal{V} such that $\partial_- \mathcal{W}' \neq \emptyset$, $\partial V \cap \mathcal{W}' = \emptyset$, and $\partial W \cap \mathcal{V}' = \emptyset$.

We call it a “type (b)- \mathcal{W} GHS” for the cases (bi) and (bii) or “type (b)- \mathcal{V} GHS” for the cases (biii) and (biv), respectively.

- (c) Each of $\partial_- \mathcal{V}_2$ and $\partial_- \mathcal{W}_1$ is disconnected but $\partial_- \mathcal{V}_2 \cap \partial_- \mathcal{W}_1$ is connected, where V and W cut off compression bodies \mathcal{V}' and \mathcal{W}' of genus at least one from \mathcal{V} and \mathcal{W} , respectively, such that $\partial_- \mathcal{V}' \neq \emptyset$, $\partial_- \mathcal{W}' \neq \emptyset$, $\partial V \cap \mathcal{W}' = \emptyset$, and $\partial W \cap \mathcal{V}' = \emptyset$. We call it a “type (c) GHS”.
- (d) Each of $\partial_- \mathcal{V}_2$ and $\partial_- \mathcal{W}_1$ consists of two components and $\partial_- \mathcal{V}_2 = \partial_- \mathcal{W}_1$, where V and W are nonseparating in \mathcal{V} and \mathcal{W} , respectively, and $\partial V \cup \partial W$ is separating in F . We call it a “type (d) GHS”.

Note that \bar{F}_{VW} is disconnected only for type (d) GHS and connected otherwise. This means if $\partial_- \mathcal{V}_2$ (resp. $\partial_- \mathcal{W}_1$) is disconnected, then $\partial_- \mathcal{V}_2 \cap \partial M \neq \emptyset$ (resp. $\partial_- \mathcal{W}_1 \cap \partial M \neq \emptyset$), excluding the case (d).

Proof. Recall that F_{VW} cannot have a S^2 -component by Lemma 2.3. Therefore, ∂W (resp. ∂V) is essential in the relevant component of F_V (resp. F_W).

Let \mathbf{H}' be the GHS obtained by preweak reduction along (V, W) from $(\mathcal{V}, \mathcal{W}; F)$. We will denote $\text{Thick}(\mathbf{H}') \cap \mathcal{V}$ and $\text{Thick}(\mathbf{H}') \cap \mathcal{W}$ as \tilde{F}_V and \tilde{F}_W , respectively, and $\text{Thin}(\mathbf{H}')$ as F_{VW} .

Since F_{VW} consists of at most three components, we divide the proof into three cases according to the number of components. In each case, we will (i) do the preweak reduction along (V, W) and then clean each product region in \mathbf{H}' , (ii) considering the remaining components of $\text{Thick}(\mathbf{H}')$ and $\overline{\text{Thin}(\mathbf{H}')}$ in \mathbf{H}' after cleaning, confirm the thick levels \tilde{F}_V and \tilde{F}_W and the inner thin level \bar{F}_{VW} , and (iii) finally determine the shapes of the region between \tilde{F}_V and \bar{F}_{VW} which would be \mathcal{V}_2 and that between \tilde{F}_W and \bar{F}_{VW} which would be \mathcal{W}_1 . Note that the idea of (iii) is inspired by the proof of Theorem 3.1 of [9].

Case a. F_{VW} consists of one component.

In this case, V and W are nonseparating in the relevant compression bodies and $\partial V \cup \partial W$ is also nonseparating in F . Hence, each of \tilde{F}_V and \tilde{F}_W consists of a genus $(n-1)$ surface and F_{VW} consists of a genus $(n-2)$ surface. Therefore, there is no product region in \mathbf{H}' to clean, i.e. $\bar{F}_V = \tilde{F}_V$, $\bar{F}_W = \tilde{F}_W$, $\bar{F}_{VW} = F_{VW}$ and they divide M into four compression bodies \mathcal{V}_1 , \mathcal{V}_2 , \mathcal{W}_1 and \mathcal{W}_2 with respect to the order of the surfaces.




Fig.2. The five types of GHSs obtained by weak reductions

Considering \mathcal{V}_2 , it is the union of the product region $\tilde{\mathcal{V}}$ between F_V and \bar{F}_V in \mathcal{V} and the 2-handle corresponding to a product neighborhood of W in \mathcal{W} , say $N_{\mathcal{W}}(W)$, where $N_{\mathcal{W}}(W) \cap \tilde{\mathcal{V}}$ is a nonseparating annulus in F_V by the assumption that $\partial V \cup \partial W$ is nonseparating in F , i.e. $\partial_- \mathcal{V}_2$ is connected. Similarly, we can use the symmetric argument for \mathcal{W}_1 and therefore $\partial_- \mathcal{W}_1$ is also connected. This means $\partial_- \mathcal{V}_2 = \partial_- \mathcal{W}_1 = \bar{F}_{VW}$. Therefore, we reach (ai) (see the first one of Figure 8 of [5]).

Case b. F_{VW} consists of two components.

Case b-i. V is separating in \mathcal{V} and W is nonseparating in \mathcal{W} .

Let F'_V be the component of F_V containing ∂W and F''_V the other component. Then we can assume F''_V is a component of $\overline{\text{Thin}}(\mathbf{H}') (= F_{VW})$. Let \tilde{F}'_V and \tilde{F}''_V be the components of $\text{Thick}(\mathbf{H}')$ obtained by pushing F'_V and F''_V off into $\text{int}(\mathcal{V})$, respectively, and let $m (< n)$ be the genus of F'_V . Since ∂W is essential in F'_V and F_{VW} cannot have a S^2 component, we get $m \geq 2$.

Here, \tilde{F}''_V and F''_V cobound a product region and therefore \tilde{F}''_V and F''_V disappear after cleaning the relevant portion of \mathbf{H}' . Hence, only \tilde{F}'_V remains in $\text{Thick}(\mathbf{H}') \cap \mathcal{V}$.

Considering Lemma 2.14, ∂W is nonseparating in F'_V and therefore $(F'_V)_W$ is a connected surface of genus $(m - 1)$, i.e. only one component $(F'_V)_W$ remains in $\overline{\text{Thin}}(\mathbf{H}')$ at this time.

Moreover, \tilde{F}_W consists of a genus $(n - 1)$ surface because W is nonseparating in \mathcal{W} . Therefore, considering $(m - 1) < (n - 1)$, the region between $(F'_V)_W$ and \tilde{F}_W cannot be homeomorphic to a product. Similarly, considering $(m - 1) < m$, the region between $(F'_V)_W$ and \tilde{F}'_V cannot be homeomorphic to a product. Hence, there is no product region to clean \mathbf{H}' at this moment.

Therefore, we get $\bar{F}_V = \tilde{F}'_V$, $\bar{F}_{VW} = (F'_V)_W$, and $\bar{F}_W = \tilde{F}_W$ and they divide M into four compression bodies \mathcal{V}_1 , \mathcal{V}_2 , \mathcal{W}_1 and \mathcal{W}_2 with respect to the order of the surfaces.

Considering \mathcal{V}_2 , it is the union of the product region $\tilde{\mathcal{V}}$ between F'_V and \tilde{F}'_V in \mathcal{V} and the 2-handle corresponding to a product neighborhood of W in \mathcal{W} , say $N_{\mathcal{W}}(W)$, where $N_{\mathcal{W}}(W) \cap \tilde{\mathcal{V}}$ is a nonseparating annulus in F'_V as we saw previously, and therefore $\partial_- \mathcal{V}_2$ is connected.

Case b-i-1. V cuts off a handlebody \mathcal{V}' from \mathcal{V} such that $\partial W \cap \mathcal{V}' = \emptyset$.

Let us expand \mathcal{V}' in \mathcal{V} until \mathcal{V}' intersects $\tilde{F}_{VW} = (F'_V)_W$ in the scar of V and let \mathcal{V}'_* be the resulting one. Then we can see \mathcal{W}_1 is the union of the product region $\tilde{\mathcal{W}}$ between F_W and \tilde{F}_W in \mathcal{W} and \mathcal{V}'_* along a once-punctured surface in F and therefore \mathcal{V}'_* does not add another component to $\partial_- \mathcal{W}_1$ other than $(F'_V)_W$ itself, i.e. $\partial_- \mathcal{W}_1$ is connected. Therefore, we reach (aii) (see the second one of Figure 8 of [5]).

Case b-i-2. V cuts off a compression body \mathcal{V}' from \mathcal{V} such that $\partial_- \mathcal{V}' \neq \emptyset$ and $\partial W \cap \mathcal{V}' = \emptyset$.

Using the similar argument as in Case b-i-1, we can see $\partial_- \mathcal{W}_1$ is disconnected because \mathcal{V}'_* adds another component to $\partial_- \mathcal{W}_1$ other than $(F'_V)_W$. Therefore, we reach (bi). Moreover, $\partial_- \mathcal{W}_1 \cap \partial M \subset \partial_- \mathcal{V}$ (see the first one of Figure 9 of [5]).

Case b-ii. V is nonseparating in \mathcal{V} and W is separating in \mathcal{W} .

In this case, we use the symmetric argument of Case b-i and therefore \tilde{F}_{VW} is connected and $\partial_- \mathcal{W}_1$ is connected.

Case b-ii-1. W cuts off a handlebody \mathcal{W}' from \mathcal{W} such that $\partial V \cap \mathcal{W}' = \emptyset$.

In this case, $\partial_- \mathcal{V}_2$ is connected and therefore we reach (aiii) by using the symmetric argument of Case b-i-1.

Case b-ii-2. W cuts off a compression body \mathcal{W}' such that $\partial_- \mathcal{W}' \neq \emptyset$ and $\partial V \cap \mathcal{W}' = \emptyset$.

In this case, $\partial_- \mathcal{V}_2$ is disconnected and therefore we reach (biii) by using the symmetric argument of Case b-i-2. Moreover, $\partial_- \mathcal{V}_2 \cap \partial M \subset \partial_- \mathcal{W}$.

Case b-iii. V and W are nonseparating in \mathcal{V} and \mathcal{W} , respectively, and $\partial V \cup \partial W$ is separating in F .

In this case, each of F_V and F_W consists of a genus $(n-1)$ surface and F_{VW} is disconnected, where each component of F_{VW} is of genus at most $(n-2)$ because ∂V (resp. ∂W) is an essential separating curve in the genus $(n-1)$ surface F_W (resp. F_V) by the assumption that $\partial V \cup \partial W$ is separating in F and there is no S^2 component of F_{VW} . Therefore, we cannot find a product region in \mathbf{H}' to clean, i.e. $\tilde{F}_V = \tilde{F}'_V$, $\tilde{F}_W = \tilde{F}'_W$, $\tilde{F}_{VW} = F_{VW}$ and they divide M into four compression bodies \mathcal{V}_1 , \mathcal{V}_2 , \mathcal{W}_1 and \mathcal{W}_2 with respect to the order of the surfaces.

Let us consider \mathcal{V}_2 . Then it is the union of the product region $\tilde{\mathcal{V}}$ between F_V and \tilde{F}_V in \mathcal{V} and the 2-handle corresponding to a product neighborhood of W in \mathcal{W} , say $N_{\mathcal{W}}(W)$, where $N_{\mathcal{W}}(W) \cap \tilde{\mathcal{V}}$ is a separating annulus in F_V . Hence, $\partial_- \mathcal{V}_2$ consists of exactly two components and therefore $\partial_- \mathcal{V}_2$ is \tilde{F}_{VW} itself. By the symmetric argument, $\partial_- \mathcal{W}_1 = \tilde{F}_{VW}$. Hence, we reach (d) (see Figure 11 of [5]).

Case c. F_{VW} consists of three components.

If both V and W are nonseparating, then F_{VW} consists of at most two components. If one

of V and W is nonseparating and the other is separating, say W is nonseparating, then ∂W must be separating in the component of F_V containing ∂W because F_{VW} consists of three components, violating Lemma 2.14.

Therefore, V and W are separating in \mathcal{V} and \mathcal{W} , respectively. Let F'_V (resp. F'_W) be the component of F_V (resp. F_W) containing ∂W (resp. ∂V) and F''_V (resp. F''_W) the other component, i.e. we can assume F''_V (resp. F''_W) is a component of $\overline{\text{Thin}(\mathbf{H})}$ ($= F_{VW}$). Let $m_{F'_V}$ ($< n$) (resp. $m_{F'_W}$ ($< n$)) be the genus of F'_V (resp. F'_W). Then we can see the following:

- (1) F_{VW} consists of three components F''_V , F''_W and F''_{VW} , where F''_{VW} is the component of F_{VW} having scars of both V and W and F''_V (resp. F''_W) has only one scar of V (resp. W). Here, the genus of F''_{VW} is

$$n - g(F''_V) - g(F''_W) = n - (n - m_{F'_V}) - (n - m_{F'_W}) = m_{F'_V} + m_{F'_W} - n.$$

- (2) \tilde{F}_V (resp. \tilde{F}_W) consists of two components \tilde{F}'_V and \tilde{F}''_V (resp. \tilde{F}'_W and \tilde{F}''_W) obtained by pushing F'_V and F''_V off into $\text{int}(\mathcal{V})$ (resp. F'_W and F''_W off into $\text{int}(\mathcal{W})$), respectively.

Then \tilde{F}''_V and F''_{VW} cobound a product region in \mathcal{V} and therefore \tilde{F}''_V and F''_{VW} disappear after cleaning the relevant portion of \mathbf{H}' . Similarly, \tilde{F}''_W and F''_{VW} disappear after cleaning the relevant portion of \mathbf{H}' . Hence, only F''_{VW} remains in $\overline{\text{Thin}(\mathbf{H})}$ and only \tilde{F}'_V and \tilde{F}'_W remain in $\text{Thick}(\mathbf{H})$ after these steps. Considering $(m_{F'_V} + m_{F'_W} - n) - m_{F'_V} = m_{F'_W} - n < 0$, we get $g(F''_{VW}) < g(\tilde{F}'_V)$ and also we get $g(F''_{VW}) < g(\tilde{F}'_W)$ likewise. Therefore, we conclude there is no product region in \mathbf{H}' to clean at this moment, i.e. $\bar{F}_V = \tilde{F}'_V$, $\bar{F}_{VW} = F''_{VW}$, and $\bar{F}_W = \tilde{F}'_W$.

Here, we can see V and W cut off compression bodies \mathcal{V}' and \mathcal{W}' from \mathcal{V} and \mathcal{W} , respectively, such that $\partial W \cap \mathcal{V}' = \emptyset$ and $\partial V \cap \mathcal{W}' = \emptyset$.

Case c-i. \mathcal{V}' and \mathcal{W}' are handlebodies.

Let us expand \mathcal{V}' (resp. \mathcal{W}') in \mathcal{V} (resp. in \mathcal{W}) until \mathcal{V}' (resp. \mathcal{W}') intersects $\bar{F}_{VW} = F''_{VW}$ in the scar of V (resp. the scar of W) and let \mathcal{V}'_* (resp. \mathcal{W}'_*) be the resulting one. Then \mathcal{W}_1 is the union of the product region $\tilde{\mathcal{W}}$ between F'_W and \bar{F}_W and \mathcal{V}'_* along a once-punctured surface in F and therefore \mathcal{V}'_* does not add another component to $\partial_- \mathcal{W}_1$ other than \bar{F}_{VW} itself, i.e. $\partial_- \mathcal{W}_1$ is connected. The symmetric argument also holds for \mathcal{V}_2 by using \mathcal{W}'_* and therefore $\partial_- \mathcal{V}_2$ is connected. This means $\partial_- \mathcal{V}_2 = \partial_- \mathcal{W}_1 = \bar{F}_{VW}$ and this leads to (aiv) (see the last one of Figure 8 of [5]).

Case c-ii. \mathcal{V}' is a compression body such that $\partial_- \mathcal{V}' \neq \emptyset$ and \mathcal{W}' is a handlebody.

Using the similar argument of Case c-i, we can see $\partial_- \mathcal{V}_2$ is connected. But $\partial_- \mathcal{W}_1$ is disconnected because \mathcal{V}'_* adds another component to $\partial_- \mathcal{W}_1$ other than \bar{F}_{VW} . Therefore, we reach (bii). Moreover, $\partial_- \mathcal{W}_1 \cap \partial M \subset \partial_- \mathcal{V}$ (see the second one of Figure 9 of [5]).

Case c-iii. \mathcal{V}' is a handlebody and \mathcal{W}' is a compression body such that $\partial_- \mathcal{W}' \neq \emptyset$.

Using the symmetric argument of Case c-ii, we can see $\partial_- \mathcal{W}_1$ is connected and $\partial_- \mathcal{V}_2$ is disconnected. Therefore, we reach (biv). Moreover, $\partial_- \mathcal{V}_2 \cap \partial M \subset \partial_- \mathcal{W}$.

Case c-iv. \mathcal{V}' and \mathcal{W}' are compression bodies such that $\partial_- \mathcal{V}' \neq \emptyset$ and $\partial_- \mathcal{W}' \neq \emptyset$.

Using the similar argument of the previous cases, we can see both $\partial_- \mathcal{V}_2$ and $\partial_- \mathcal{W}_1$ are

disconnected. Therefore, we reach (c). Moreover, $\partial_- \mathcal{W}_1 \cap \partial M \subset \partial_- \mathcal{V}$ and $\partial_- \mathcal{V}_2 \cap \partial M \subset \partial_- \mathcal{W}$ (see Figure 10 of [5]).

This completes the proof. \square

As summary of the proof of Lemma 3.1, we get the following corollary directly.

Corollary 3.2. *Let M , F and \mathbf{H} be as in Lemma 3.1. Then the GHS $\mathbf{H} = (\mathcal{V}_1, \mathcal{V}_2; \bar{F}_V) \cup_{\bar{F}_{VW}} (\mathcal{W}_1, \mathcal{W}_2; \bar{F}_W)$ is obtained as follows.*

- (1) *The thick level \bar{F}_V (resp. \bar{F}_W) is obtained by pushing the component of F_V (resp. F_W) containing ∂W (resp. ∂V) off slightly into $\text{int}(\mathcal{V})$ (resp. into $\text{int}(\mathcal{W})$).*
- (2) *The inner thin level \bar{F}_{VW} is the union of components of F_{VW} having scars of both V and W . Moreover, if $\partial_- \mathcal{V}_2 \cap \partial M \neq \emptyset$ (resp. $\partial_- \mathcal{W}_1 \cap \partial M \neq \emptyset$), then it belongs to $\partial_- \mathcal{W}$ (resp. $\partial_- \mathcal{V}$).*

Lemma 3.3. *Let M and F be as in Lemma 3.1. If two GHSs \mathbf{H}_1 and \mathbf{H}_2 obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$ are isotopic in M , then they are of the same type in the sense of Lemma 3.1.*

Proof. Let $\mathbf{H}_i = (\mathcal{V}_1^i, \mathcal{V}_2^i; \bar{F}_{V_i}) \cup_{\bar{F}_{V_i W_i}} (\mathcal{W}_1^i, \mathcal{W}_2^i; \bar{F}_{W_i})$, where $\mathcal{V}_2^i \cap \mathcal{W}_1^i = \bar{F}_{V_i W_i}$ for $i = 1, 2$. Suppose there is an isotopy $h_t : M \rightarrow M$, $t \in [0, 1]$ such that $h_1(\mathbf{H}_1) = \mathbf{H}_2$. Since an isotopy preserves each component of ∂M , we will not consider such components of $\text{Thin}(\mathbf{H}_1)$ and $\text{Thin}(\mathbf{H}_2)$.

If there is a collection of mutually disjoint surfaces C (assume each element of C is connected), then we can divide C into two collections C_1 and C_2 such that (i) each element of C_1 cuts off a submanifold of M intersecting only itself among the elements of C and missing the others and (ii) each element of C_2 does not. Then we will say “ C_1 is the type (i) subset of C and C_2 is the type (ii) subset of C ”. We can see the type (i) subset of C misses the type (ii) subset of C and the union of these two subsets is the entire collection C . Let \mathbf{H} be a GHS obtained by weak reduction from $(\mathcal{V}, \mathcal{W}; F)$. Putting $C = \text{Thick}(\mathbf{H}) \cup \overline{\text{Thin}(\mathbf{H})}$, each element of $\text{Thick}(\mathbf{H})$ cuts off a compression body intersecting only itself among the elements of C and each element of $\overline{\text{Thin}(\mathbf{H})}$ does not cut off such a submanifold by Lemma 3.1 (see Fig.2). This means $\text{Thick}(\mathbf{H})$ is the type (i) subset of C and $\overline{\text{Thin}(\mathbf{H})}$ is the type (ii) subset of C . Note that if there is an ambient isotopy g_t defined on M , then the type (i) and type (ii) subsets of $g_t(C)$ are the images of the type (i) and type (ii) subsets of C of g_t , respectively.

Therefore, $h_1(\text{Thick}(\mathbf{H}_1))$ and $h_1(\overline{\text{Thin}(\mathbf{H}_1)})$ are the type (i) and type (ii) subsets of $h_1(\text{Thick}(\mathbf{H}_1) \cup \overline{\text{Thin}(\mathbf{H}_1)})$, respectively. But the assumption that $h_1(\mathbf{H}_1) = \mathbf{H}_2$ implies $h_1(\text{Thick}(\mathbf{H}_1) \cup \overline{\text{Thin}(\mathbf{H}_1)}) = \text{Thick}(\mathbf{H}_2) \cup \overline{\text{Thin}(\mathbf{H}_2)}$. Since $\text{Thick}(\mathbf{H}_2)$ and $\overline{\text{Thin}(\mathbf{H}_2)}$ are the type (i) and type (ii) subsets of $\text{Thick}(\mathbf{H}_2) \cup \overline{\text{Thin}(\mathbf{H}_2)}$, respectively, we conclude $h_1(\text{Thick}(\mathbf{H}_1)) = \text{Thick}(\mathbf{H}_2)$ and $h_1(\overline{\text{Thin}(\mathbf{H}_1)}) = \overline{\text{Thin}(\mathbf{H}_2)}$.

This means $h_1(\mathcal{V}_2^1, \mathcal{W}_1^1) = (\mathcal{V}_2^2, \mathcal{W}_1^2)$ or $(\mathcal{W}_1^2, \mathcal{V}_2^2)$, i.e. at least one of $\partial_- \mathcal{V}_2^1 \cap \partial M$ and $\partial_- \mathcal{W}_1^1 \cap \partial M$ is not empty if and only if at least one of $\partial_- \mathcal{V}_2^2 \cap \partial M$ and $\partial_- \mathcal{W}_1^2 \cap \partial M$ is not empty.

Case a. $\partial_- \mathcal{V}_2^i \cap \partial M = \emptyset$ and $\partial_- \mathcal{W}_1^i \cap \partial M = \emptyset$ for $i = 1, 2$.

In this case, considering Lemma 3.1, \mathbf{H}_1 and \mathbf{H}_2 are of type (a) or type (d). Since

$h_1(\bar{F}_{V_1W_1}) = \bar{F}_{V_2W_2}$, the number of components of $\bar{F}_{V_1W_1}$ is the same as that of $\bar{F}_{V_2W_2}$. This means both \mathbf{H}_1 and \mathbf{H}_2 are either of type (a) or of type (d).

Case b. At least one of $\partial_- \mathcal{V}_2^i \cap \partial M$ and $\partial_- \mathcal{W}_1^i \cap \partial M$ is not empty for $i = 1, 2$.

In this case, \mathbf{H}_1 and \mathbf{H}_2 are of type (b)- \mathcal{W} , type (b)- \mathcal{V} or type (c).

Without loss of generality, assume $\partial_- \mathcal{V}_2^1 \cap \partial M \neq \emptyset$. In this case, $\mathcal{V}_2^1 \cap \partial M = \partial_- \mathcal{V}_2^1 \cap \partial M \subset \partial_- \mathcal{W}$ by Corollary 3.2. Recall $h_1(\mathcal{V}_2^1)$ is either \mathcal{V}_2^2 or \mathcal{W}_1^2 . In the latter case, $h_1(\mathcal{V}_2^1) \cap \partial M = \mathcal{W}_1^2 \cap \partial M \subset \partial_- \mathcal{V}$ by Corollary 3.2, leading to a contradiction because an isotopy preserves each component of ∂M . Hence, $h_1(\mathcal{V}_2^1) = \mathcal{V}_2^2$ and $h_1(\mathcal{W}_1^1) = \mathcal{W}_1^2$. Therefore, we conclude the shapes of $\partial_- \mathcal{V}_2^1$ and $\partial_- \mathcal{W}_1^1$ (including the property whether each of them intersects ∂M or not) are the same as those of $\partial_- \mathcal{V}_2^2$ and $\partial_- \mathcal{W}_1^2$, respectively. Hence, \mathbf{H}_1 and \mathbf{H}_2 are of the same type in the sense of Lemma 3.1.

This completes the proof. \square

Here, we introduce important ideas in the author's result [4] according to the context.

DEFINITION 3.4 (MODIFICATION OF DEFINITION 2.12 OF [4]). Let M and F be as in Lemma 3.1 and recall the equivalence relation \sim defined on \mathcal{GHS}_F^* in Lemma 2.7. In a weak reducing pair for the Heegaard splitting $(\mathcal{V}, \mathcal{W}; F)$, if a disk belongs to \mathcal{V} , then we call it a \mathcal{V} -disk. Otherwise, we call it a \mathcal{W} -disk. We call a 2-simplex in $D(F)$ represented by two vertices in $D_{\mathcal{V}}(F)$ and one vertex in $D_{\mathcal{W}}(F)$ a \mathcal{V} -face, and also define a \mathcal{W} -face symmetrically. If there is a \mathcal{V} - or \mathcal{W} -face Δ , then there exist exactly two weak reducing pairs in Δ and one shares a disk with the other. We say the two weak reducing pairs *form* Δ . Suppose there is a \mathcal{V} -face Δ such that the GHSs obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$ along the two weak reducing pairs forming Δ are equivalent. Then we call this \mathcal{V} -face a “ \mathcal{V} -face having one equivalent class (we will use *OEC* as abbreviation from now on)” and also we define a “ \mathcal{W} -face having *OEC*” likewise.

Let us consider a graph as follows.

- (1) We assign a vertex to each \mathcal{V} -face having *OEC*.
- (2) If a \mathcal{V} -face having *OEC* shares a weak reducing pair with another \mathcal{V} -face having *OEC*, then we assign an edge between these two vertices in the graph.

We call this graph the *graph of \mathcal{V} -faces having *OEC**. If there is a maximal subset $\varepsilon_{\mathcal{V}}$ of \mathcal{V} -faces having *OEC* which represents a connected component of the graph of \mathcal{V} -faces having *OEC* and the component is not an isolated vertex, then we call $\varepsilon_{\mathcal{V}}$ a *\mathcal{V} -facial cluster having *OEC**. Similarly, we define the *graph of \mathcal{W} -faces having *OEC** and a *\mathcal{W} -facial cluster having *OEC**. Although each \mathcal{V} -face having *OEC* forming a \mathcal{V} -facial cluster having *OEC* induces one equivalent class, it is not obvious that the facial cluster also induces one equivalent class because the number of \mathcal{V} -faces in the cluster might be infinite. We will clarify the meaning of the phrase “*having *OEC**” in Lemma 3.7.

Lemma 3.5. *Let M and F be as in Lemma 3.1. Let Δ be a \mathcal{V} -face. Then the GHSs obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$ along the two weak reducing pairs forming Δ are equivalent if and only if one \mathcal{V} -disk of Δ cuts off a solid torus \mathcal{V}' from \mathcal{V} and the other \mathcal{V} -disk is a meridian disk of \mathcal{V}' .*

Proof. Let $\Delta = \{V_1, V_2, W\}$ for $V_1, V_2 \subset \mathcal{V}$ and $W \subset \mathcal{W}$, and $\mathbf{H}_i = (\bar{F}_{V_i}, \bar{F}_{V_i W}, \bar{F}_W^i)$ the GHS obtained by weak reduction along (V_i, W) for $i = 1, 2$.

(\Leftarrow) Suppose V_1 cuts off a solid torus \mathcal{V}' from \mathcal{V} and V_2 is a meridian disk of \mathcal{V}' (so V_2 is nonseparating in \mathcal{V}). Let F'_{V_1} be the genus $(n-1)$ -component of F_{V_1} . Since $\partial W \cap \mathcal{V}' = \emptyset$ by Lemma 2.3, F'_{V_1} contains ∂W . This means the component of F_W containing ∂V_1 , say F'_W , also contains the once-punctured torus that ∂V_1 cuts off from F . Hence, F'_W also contains ∂V_2 and therefore we can assume $\bar{F}_W^1 = \bar{F}_W^2$ by Corollary 3.2.

Here, \bar{F}_{V_1} and \bar{F}_{V_2} are obtained by pushing F'_{V_1} and F_{V_2} off slightly into $\text{int}(\mathcal{V})$, respectively, by Corollary 3.2. Moreover, considering (2.11) of Lemma 2.11 and the genus of \bar{F}_{V_2} , \bar{F}_{V_2} is isotopic to \bar{F}_{V_1} in \mathcal{V} . Indeed, considering (c) and (d) of Fig.1, we can say the isotopy taking \bar{F}_{V_2} into \bar{F}_{V_1} pushes off the 3-ball part in (c) into $\mathcal{V} - \mathcal{V}'$ without affecting a small neighborhood of V_2 and that of W in M . Hence, considering the component of $F_{V_2 W}$ having scars of both V_2 and W , it is isotopic to the component of $F_{V_1 W}$ having scars of both V_1 and W in M by the similar way sending \bar{F}_{V_2} into \bar{F}_{V_1} . This means $\bar{F}_{V_2 W}$ is isotopic to $\bar{F}_{V_1 W}$ in M by Corollary 3.2. Therefore, we can isotope \mathbf{H}_2 into \mathbf{H}_1 by isotoping \bar{F}_{V_2} into \bar{F}_{V_1} first and then isotoping $\bar{F}_{V_2 W}$ into $\bar{F}_{V_1 W}$ next.

Hence, summing up the above observations, we conclude \mathbf{H}_2 is equivalent to \mathbf{H}_1 .

(\Rightarrow) Suppose \mathbf{H}_1 is equivalent to \mathbf{H}_2 . Hence, \bar{F}_{V_1} is isotopic to \bar{F}_{V_2} in \mathcal{V} .

Case a. Both V_1 and V_2 are nonseparating in \mathcal{V} .

In this case, \bar{F}_{V_1} and \bar{F}_{V_2} are obtained by pushing F_{V_1} and F_{V_2} off slightly into $\text{int}(\mathcal{V})$, respectively, by Corollary 3.2. Therefore, considering the assumption that \bar{F}_{V_1} is isotopic to \bar{F}_{V_2} in \mathcal{V} , V_1 is isotopic to V_2 in \mathcal{V} by (2.11) of Lemma 2.11, violating the assumption that Δ is a 2-simplex.

Case b. One of V_1 and V_2 is nonseparating and the other is separating in \mathcal{V} .

Assume V_1 is nonseparating. In this case, \bar{F}_{V_1} is obtained by pushing F_{V_1} off slightly into $\text{int}(\mathcal{V})$ and \bar{F}_{V_2} is obtained by pushing the component of F_{V_2} containing ∂W off slightly into $\text{int}(\mathcal{V})$ by Corollary 3.2. Since V_1 is nonseparating and V_2 is separating in \mathcal{V} , the assumption that \bar{F}_{V_1} is isotopic to \bar{F}_{V_2} in \mathcal{V} induces that V_2 cuts off a solid torus \mathcal{V}' from \mathcal{V} and V_1 is isotopic to a meridian disk D of \mathcal{V}' in \mathcal{V} by (2.11) of Lemma 2.11 and we can assume that $D \cap V_2 = \emptyset$.

Considering $V_1 \cap V_2 = \emptyset$, either $V_1 \subset \mathcal{V}'$ or $V_1 \cap \mathcal{V}' = \emptyset$. Assume that $V_1 \cap \mathcal{V}' = \emptyset$, i.e. the component of $F - \partial V_2$ containing ∂D is different from that containing ∂V_1 . Then ∂V_1 and ∂D bound an annulus A in F because they are disjoint, isotopic, essential simple closed curves in F (see Lemma 3.3 of [10]). If $A \cap \partial V_2 = \emptyset$, then both ∂V_1 and ∂D belong to one component of $F - \partial V_2$, leading to a contradiction. Therefore, considering $(D \cup V_1) \cap V_2 = \emptyset$, $\partial V_2 \subset A$ and it is essential in A because V_2 is a compressing disk for F . This means ∂V_2 is isotopic to ∂V_1 in A and therefore V_2 is nonseparating in \mathcal{V} , violating the assumption.

Therefore, $V_1 \subset \mathcal{V}'$. Hence, considering Lemma 2.14, V_1 is a meridian disk of \mathcal{V}' .

Case c. Both V_1 and V_2 are separating in \mathcal{V} .

In this case, \bar{F}_{V_1} and \bar{F}_{V_2} are obtained by pushing the components of F_{V_1} and F_{V_2} con-

taining ∂W off slightly into $\text{int}(\mathcal{V})$, respectively, by Corollary 3.2. Moreover, V_1 and V_2 are mutually disjoint separating disks in \mathcal{V} . Therefore, the assumption that \bar{F}_{V_1} is isotopic to \bar{F}_{V_2} in \mathcal{V} induces V_1 is isotopic to V_2 in \mathcal{V} by Lemma 2.13. This violates the assumption that Δ is a 2-simplex.

Therefore, we can see only Case b holds, leading to the result.

This completes the proof of Lemma 3.5. \square

Considering Lemma 2.9 of [3], we get the following corollary directly from Lemma 3.5.

Corollary 3.6. *Suppose M is an irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ is a weakly reducible, unstabilized Heegaard splitting of M of genus three. Then every \mathcal{V} - or \mathcal{W} -face has OEC.*

Lemma 3.7 (analogue to Lemma 2.13 of [4]). *Let M and F be as in Lemma 3.1. If there are two different \mathcal{V} -faces having OEC $\Delta_1 = \{V_0, V_1, W\}$ and $\Delta_2 = \{V_1, V_2, W\}$, then V_1 is nonseparating and V_0 and V_2 are separating in \mathcal{V} . Therefore, there is a unique weak reducing pair in a \mathcal{V} -facial cluster having OEC which can belong to two or more faces in the \mathcal{V} -facial cluster having OEC. This means every GHS obtained by weak reduction in a \mathcal{V} -facial cluster having OEC is equivalent to the GHS obtained by using the commonly shared weak reducing pair.*

Proof. For the sake of contradiction, assume V_1 is separating in F . Then V_0 and V_2 are meridian disks of the solid torus \mathcal{V}' that V_1 cuts off from \mathcal{V} by Lemma 3.5, where this solid torus is uniquely determined by the assumption that the genus of F is at least three. But the uniqueness of a meridian disk of a solid torus up to isotopy forces Δ_1 and Δ_2 to be the same, violating the assumption. Hence, V_1 is nonseparating and therefore V_0 and V_2 are separating by Lemma 3.5. This completes the proof. \square

Considering Lemma 3.7, we can say a \mathcal{V} - or \mathcal{W} -facial cluster having OEC ε has one equivalent class in a real sense.

DEFINITION 3.8 (ANALOGUE TO DEFINITION 2.14 OF [4]). *Let M and F be as in Lemma 3.1. Considering Lemma 3.7, there is a unique weak reducing pair in a \mathcal{V} -facial cluster having OEC which can belong to two or more faces in the cluster. We call it the *center*. Lemma 3.5 and Lemma 3.7 induce any \mathcal{V} -disk in the cluster not belonging to the center cuts off a solid torus from \mathcal{V} , where this solid torus is uniquely determined by the disk, and the \mathcal{V} -disk in the center is a meridian disk of the solid torus. The shape of a \mathcal{V} -facial cluster having OEC is exactly the same as a \mathcal{V} -facial cluster which was introduced by the author for the genus three case in [4] (see the third one of Fig.4). Therefore, if a \mathcal{V} -face in a \mathcal{V} -facial cluster having OEC is represented by two weak reducing pairs, then one weak reducing pair is the center.*

Lemma 3.9 (analogue to Lemma 2.15 of [4]). *Let M and F be as in Lemma 3.1. Every \mathcal{V} -face having OEC belongs to a uniquely determined \mathcal{V} -facial cluster having OEC. Moreover, every \mathcal{V} -facial cluster having OEC has infinitely many weak reducing pairs.*

Proof. A \mathcal{V} -face Δ has OEC if and only if a \mathcal{V} -disk of Δ cuts off a solid torus \mathcal{V}' from

\mathcal{V} and the other \mathcal{V} -disk is a meridian disk of \mathcal{V}' by Lemma 3.5. Therefore, the proof is essentially the same as that of Lemma 2.15 of [4] except for the uniqueness of the \mathcal{V} -facial cluster having OEC which contains Δ .

Let us consider the uniqueness. Suppose a \mathcal{V} -face Δ having OEC belongs to two \mathcal{V} -facial clusters having OEC, say ε_1 and ε_2 . Then there is a weak reducing pair in Δ containing a nonseparating \mathcal{V} -disk by Lemma 3.5 and it must be the center of ε_i for $i = 1, 2$ by Definition 3.8. This means all \mathcal{V} -faces of ε_1 and ε_2 share the common center, i.e. ε_1 and ε_2 correspond to the same connected component of the graph of \mathcal{V} -faces having OEC and therefore we conclude $\varepsilon_1 = \varepsilon_2$.

This completes the proof. \square

The proof of Lemma 3.9 induces the following corollary directly.

Corollary 3.10. *Let M and F be as in Lemma 3.1. For a weak reducing pair, there is at most one \mathcal{V} -facial cluster having OEC whose center is the weak reducing pair.*

Lemma 3.11. *Let M and F be as in Lemma 3.1. For a \mathcal{V} -facial cluster having OEC, say $\varepsilon_{\mathcal{V}}$, if V_1 and V_2 are different \mathcal{V} -disks in $\varepsilon_{\mathcal{V}}$ not contained in the center, then there is no 1-simplex in $D(F)$ between them.*

Proof. Let (\bar{V}, \bar{W}) be the center of $\varepsilon_{\mathcal{V}}$. For the sake of contradiction, assume that there is a 1-simplex between V_1 and V_2 in $D(F)$. Considering Definition 3.8, ∂V_1 and ∂V_2 cut off once-punctured tori T_1 and T_2 from F , respectively, where T_i is the closure of the component of $F - \partial V_i$ containing $\partial \bar{V}$ for $i = 1, 2$. Moreover, considering the assumption that $\partial V_1 \cap \partial V_2 = \emptyset$, one of T_1 and T_2 contains the other, say $T_1 \subset T_2$. Here, ∂V_1 divides T_2 into two pieces, where one is a genus one surface and the other is a planar surface. Since V_1 is a compressing disk of F , ∂V_1 cannot cut off a disk from T_2 . This means ∂V_1 cuts off an annulus from T_2 such that the other boundary component is ∂V_2 , i.e. ∂V_1 is isotopic to ∂V_2 in F . Therefore, V_1 is isotopic to V_2 in \mathcal{V} by Lemma 2.1, leading to a contradiction. This completes the proof. \square

Corollary 3.12. *Let M and F be as in Lemma 3.1. For a \mathcal{V} -facial cluster having OEC, say $\varepsilon_{\mathcal{V}}$, there is no n -simplex in $D(F)$ spanned by vertices of $\varepsilon_{\mathcal{V}}$ for $n \geq 3$.*

Proof. Let (\bar{V}, \bar{W}) be the center of $\varepsilon_{\mathcal{V}}$. For the sake of contradiction, assume that there is an n -simplex Δ in $D(F)$ spanned by vertices of $\varepsilon_{\mathcal{V}}$ for $n \geq 3$. Then we can obtain an at least $(n - 2)$ -subsimplex δ by removing \bar{V} and \bar{W} from Δ , where $n - 2 \geq 1$. Here, δ consists of \mathcal{V} -disks of $\varepsilon_{\mathcal{V}}$ other than \bar{V} , violating Lemma 3.11. \square

Lemma 3.13. *Let M and F be as in Lemma 3.1. For a \mathcal{V} -facial cluster having OEC, say $\varepsilon_{\mathcal{V}}$, the union of all simplices of $D(F)$ spanned by the vertices of $\varepsilon_{\mathcal{V}}$ is $\varepsilon_{\mathcal{V}}$ itself.*

Proof. Considering that each \mathcal{V} -face of $\varepsilon_{\mathcal{V}}$ is spanned by three vertices of $\varepsilon_{\mathcal{V}}$, $\varepsilon_{\mathcal{V}}$ is contained in the union of all simplices of $D(F)$ spanned by the vertices of $\varepsilon_{\mathcal{V}}$. Hence, we will prove that the union belongs to $\varepsilon_{\mathcal{V}}$. It is sufficient to show that every simplex of $D(F)$ spanned by the vertices of $\varepsilon_{\mathcal{V}}$ is contained in $\varepsilon_{\mathcal{V}}$.

Let (\bar{V}, \bar{W}) be the center of $\varepsilon_{\mathcal{V}}$. Suppose there is a 1-simplex σ spanned by two vertices of $\varepsilon_{\mathcal{V}}$ which is not contained in $\varepsilon_{\mathcal{V}}$. If there is a vertex of $\varepsilon_{\mathcal{V}}$, then it is either (i) \bar{W} , (ii) \bar{V} , or

(iii) a \mathcal{V} -disk other than \bar{V} . Considering all combinations of two different vertices of $\varepsilon_{\mathcal{V}}$ and the shape of $\varepsilon_{\mathcal{V}}$ (see the third one of Fig.4), if they are of different types among the previous three types, then $\sigma \subset \varepsilon_{\mathcal{V}}$, violating the assumption. This means σ must connect two \mathcal{V} -disks other than \bar{V} , violating Lemma 3.11. Hence, every 1-simplex spanned by two vertices of $\varepsilon_{\mathcal{V}}$ must belong to $\varepsilon_{\mathcal{V}}$.

Suppose there is a 2-simplex Δ spanned by three vertices of $\varepsilon_{\mathcal{V}}$ which is not contained in $\varepsilon_{\mathcal{V}}$. Considering all combinations of three vertices of Δ similarly as in the previous paragraph, if three vertices are of mutually different types, then Δ belongs to $\varepsilon_{\mathcal{V}}$, violating the assumption of Δ . Hence, there must be two \mathcal{V} -disks other than \bar{V} in Δ , violating Lemma 3.11. Therefore, every 2-simplex spanned by three vertices of $\varepsilon_{\mathcal{V}}$ must belong to $\varepsilon_{\mathcal{V}}$.

Since there is no simplex of dimension at least three spanned by vertices of $\varepsilon_{\mathcal{V}}$ by Corollary 3.12, we do not need to consider such simplices.

This completes the proof. \square

From now on, we will find a collection of connected subsets of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ such that (i) the GHSs obtained by weak reductions along the weak reducing pairs in such a subset are all equivalent and (ii) every weak reducing pair belongs to exactly one subset among such subsets (see Lemma 3.29) in Definition 3.19, Definition 3.22, Definition 3.23 and Definition 3.24.

We will subdivide each of the five types of GHSs obtained by weak reductions in Lemma 3.1 as in the following definition.

DEFINITION 3.14. Let M and F be as in Lemma 3.1. Let \mathbf{H} be the GHS obtained by weak reduction along a weak reducing pair (V, W) from $(\mathcal{V}, \mathcal{W}; F)$. We subdivide each of the five types of GHSs of Lemma 3.1, where we will differentiate (1) the case where V or W is nonseparating or it cuts off a solid torus from the relevant compression body and (2) the other cases in each subcase. Note that if V cuts off a solid torus \mathcal{V}' from \mathcal{V} , then \mathcal{V}' must miss ∂W by Lemma 2.3 and the similar argument also holds for W .

(a) Case: \mathbf{H} is of type (a).

(i) If

(I) both V and W are nonseparating or

(II) if there is a separating one, then it cuts off a solid torus from the relevant compression body,

then we say \mathbf{H} is of *type (a)-(i)*.

(ii) (I) One of V and W is nonseparating or it cuts off a solid torus from the relevant compression body and (II) the other is separating and it does not cut off a solid torus from the relevant compression body.

(\mathcal{W}) If V is separating and it does not cut off a solid torus from \mathcal{V} (V cuts off a handlebody \mathcal{V}' of genus at least two from \mathcal{V} such that $\partial W \cap \mathcal{V}' = \emptyset$), then we say \mathbf{H} is of *type (a)-(ii)- \mathcal{W}* .

(\mathcal{V}) If W is separating and it does not cut off a solid torus from \mathcal{W} (W cuts off a handlebody \mathcal{W}' of genus at least two from \mathcal{W} such that $\partial V \cap \mathcal{W}' = \emptyset$), then we say \mathbf{H} is of *type (a)-(ii)- \mathcal{V}* .

(iii) If (I) both V and W are separating and (II) each of them does not cut off a solid torus from the relevant compression body (they cut off handlebodies \mathcal{V}' and \mathcal{W}'

of genera at least two from \mathcal{V} and \mathcal{W} , respectively, such that $\partial W \cap \mathcal{V}' = \emptyset$ and $\partial V \cap \mathcal{W}' = \emptyset$), then we say \mathbf{H} is of type (a)-(iii).

(b) Case: \mathbf{H} is of type (b)- \mathcal{W} or type (b)- \mathcal{V} .

(\mathcal{W}) Case: \mathbf{H} is of type (b)- \mathcal{W} (V cuts off a compression body \mathcal{V}' of genus at least one such that $\partial_- \mathcal{V}' \neq \emptyset$ and $\partial W \cap \mathcal{V}' = \emptyset$).

If W is nonseparating or it cuts off a solid torus from \mathcal{W} , then we say \mathbf{H} is of type (b)- \mathcal{W} -(i). Otherwise (W cuts off a handlebody \mathcal{W}' of genus at least two from \mathcal{W} such that $\partial V \cap \mathcal{W}' = \emptyset$), we say \mathbf{H} is of type (b)- \mathcal{W} -(ii).

(\mathcal{V}) Case: \mathbf{H} is of type (b)- \mathcal{V} (W cuts off a compression body \mathcal{W}' of genus at least one such that $\partial_- \mathcal{W}' \neq \emptyset$ and $\partial V \cap \mathcal{W}' = \emptyset$).

If V is nonseparating or it cuts off a solid torus from \mathcal{V} , then we say \mathbf{H} is of type (b)- \mathcal{V} -(i). Otherwise (V cuts off a handlebody \mathcal{V}' of genus at least two from \mathcal{V} such that $\partial W \cap \mathcal{V}' = \emptyset$), we say \mathbf{H} is of type (b)- \mathcal{V} -(ii).

(c) Case: \mathbf{H} is of type (c).

We do not subdivide this case.

(d) Case: \mathbf{H} is of type (d).

We do not subdivide this case.

For a given weak reducing pair (V, W) , we can see the GHS obtained by weak reduction along (V, W) belongs to exactly one subcase among the ten types of Definition 3.14.

Lemma 3.15. *If two GHSs $\mathbf{H}_1 = (\bar{F}_{V_1}, \bar{F}_{V_1 W_1}, \bar{F}_{W_1})$ and $\mathbf{H}_2 = (\bar{F}_{V_2}, \bar{F}_{V_2 W_2}, \bar{F}_{W_2})$ obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$ are equivalent, then they are of the same type in the sense of Definition 3.14. This induces that the GHSs obtained by weak reductions along the weak reducing pairs in a given \mathcal{V} - or \mathcal{W} -facial cluster having OEC are of the same type in the sense of Definition 3.14.*

Proof. Since \mathbf{H}_1 is isotopic to \mathbf{H}_2 in M , $\bar{F}_{V_1 W_1}$ is isotopic to $\bar{F}_{V_2 W_2}$ in M as in the proof of Lemma 3.3. This means if one of \mathbf{H}_1 and \mathbf{H}_2 is of type (d), where this is the only case where the inner thin level is disconnected, then the other is also of type (d), leading to the result. Hence, assume both \mathbf{H}_1 and \mathbf{H}_2 are not of type (d), i.e. both $\bar{F}_{V_1 W_1}$ and $\bar{F}_{V_2 W_2}$ are connected.

- CLAIM 1.** (i) V_1 is nonseparating or cuts off a solid torus missing ∂W_1 from \mathcal{V} if and only if V_2 is nonseparating or cuts off a solid torus missing ∂W_2 from \mathcal{V} .
(ii) V_1 cuts off a handlebody of genus at least two missing ∂W_1 from \mathcal{V} if and only if V_2 cuts off a handlebody of genus at least two missing ∂W_2 from \mathcal{V} .
(iii) V_1 cuts off a compression body with nonempty negative boundary missing ∂W_1 from \mathcal{V} if and only if V_2 cuts off a compression body with nonempty negative boundary missing ∂W_2 from \mathcal{V} .

Proof of Claim 1. Note that V_1 is either (i)-(1) nonseparating, (i)-(2) cuts off a solid torus missing ∂W_1 from \mathcal{V} , (ii) cuts off a handlebody of genus at least two missing ∂W_1 from \mathcal{V} , or (iii) cuts off a compression body with nonempty negative boundary missing ∂W_1 from \mathcal{V} (the same argument also holds for V_2).

Case a. At least one of V_1 and V_2 is nonseparating in \mathcal{V} .

Assume V_i is nonseparating in \mathcal{V} for $i \neq j$. If V_j is also nonseparating, then it implies a

subcase of (i) and therefore there is nothing to prove. Hence, assume that V_j is separating in \mathcal{V} . Since \bar{F}_{V_j} is isotopic to \bar{F}_{V_i} in \mathcal{V} by the assumption that \mathbf{H}_j is equivalent to \mathbf{H}_i , Corollary 3.2 and (2.11) of Lemma 2.11 induce V_j cuts off a solid torus \mathcal{V}' from \mathcal{V} and \mathcal{V}' misses ∂W_j by Lemma 2.3. This means that we have proved (i) of Claim 1 except the case where both V_1 and V_2 are separating.

This completes the proof of Case a.

Case b. Both V_1 and V_2 are separating in \mathcal{V} .

By Corollary 3.2, \bar{F}_{V_i} is obtained by pushing the component of F_{V_i} containing ∂W_i slightly into $\text{int}(\mathcal{V})$ for $i = 1, 2$. Therefore, there is a region cobounded by V_i , \bar{F}_{V_i} and a component of $F - \partial V_i$ in \mathcal{V} , say $\tilde{\mathcal{V}}_i$, which is homeomorphic to $\bar{F}_{V_i} \times I$ such that the 0-level is \bar{F}_{V_i} and the 1-level contains ∂W_i for $i = 1, 2$. Let $\tilde{\mathcal{V}}'_1$ be the closure of the component of $\mathcal{V} - \bar{F}_{V_1}$ missing $\partial_+ \mathcal{V}$ and $\tilde{\mathcal{V}}''_1$ the closure of the component of $\mathcal{V} - \tilde{\mathcal{V}}_1$ intersecting $\partial_+ \mathcal{V}$. Since \mathbf{H}_1 and \mathbf{H}_2 are equivalent, there is an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that $h_1(\bar{F}_{V_2}) = \bar{F}_{V_1}$ and therefore $h_1(\tilde{\mathcal{V}}_2)$ is homeomorphic to $\bar{F}_{V_1} \times I$ whose 0-level is \bar{F}_{V_1} . Here, $\text{cl}(\mathcal{V} - h_1(\tilde{\mathcal{V}}_2))$ consists of two components, where the component missing $\partial_+ \mathcal{V}$ is $\tilde{\mathcal{V}}'_1$ and the other component, say $\tilde{\mathcal{V}}''_2$, intersects $\partial_+ \mathcal{V}$.

Then we can see

$$\mathcal{V} = \tilde{\mathcal{V}}''_1 \cup_{V_1} (\tilde{\mathcal{V}}_1 \cup_{\bar{F}_{V_1}} \tilde{\mathcal{V}}'_1) = \tilde{\mathcal{V}}''_2 \cup_{h_1(V_2)} (h_1(\tilde{\mathcal{V}}_2) \cup_{\bar{F}_{V_1}} \tilde{\mathcal{V}}'_1),$$

where $\partial W_1 \cap \tilde{\mathcal{V}}''_1 = \emptyset$ and $h_1(\partial W_2) \cap \tilde{\mathcal{V}}''_2 = \emptyset$, and both $\tilde{\mathcal{V}}''_1$ and $\tilde{\mathcal{V}}''_2$ are compression bodies such that $\partial_- \tilde{\mathcal{V}}''_1 \subset \partial_- \mathcal{V}$ and $\partial_- \tilde{\mathcal{V}}''_2 \subset \partial_- \mathcal{V}$ by Lemma 1.3 of [12] because we get $\tilde{\mathcal{V}}''_1$ and $\tilde{\mathcal{V}}''_2$ by cutting \mathcal{V} off along V_1 and $h_1(V_2)$, respectively. Moreover, $\tilde{\mathcal{V}}_1 \cup_{\bar{F}_{V_1}} \tilde{\mathcal{V}}'_1$ and $h_1(\tilde{\mathcal{V}}_2) \cup_{\bar{F}_{V_1}} \tilde{\mathcal{V}}'_1$ are compression bodies such that one is homeomorphic to the other and they share the common negative boundary. This means (1) the genus of $\tilde{\mathcal{V}}''_1$ is the same as that of $\tilde{\mathcal{V}}''_2$ and (2) $\tilde{\mathcal{V}}''_1$ is a handlebody if and only if $\tilde{\mathcal{V}}''_2$ is a handlebody. That is, $\tilde{\mathcal{V}}''_1$ is

- (i) a solid torus (= V_1 cuts off a solid torus missing ∂W_1 from \mathcal{V}),
- (ii) a handlebody of genus at least two (= V_1 cuts off a handlebody of genus at least two missing ∂W_1 from \mathcal{V}), or
- (iii) a compression body with nonempty negative boundary (= V_1 cuts off a compression body with nonempty negative boundary missing ∂W_1 from \mathcal{V})

if and only if $\tilde{\mathcal{V}}''_2$ is

- (i) a solid torus (= $h_1(V_2)$ cuts off a solid torus missing $h_1(\partial W_2)$ from \mathcal{V}),
- (ii) a handlebody of genus at least two (= $h_1(V_2)$ cuts off a handlebody of genus at least two missing $h_1(\partial W_2)$ from \mathcal{V}), or
- (iii) a compression body with nonempty negative boundary (= $h_1(V_2)$ cuts off a compression body with nonempty negative boundary missing $h_1(\partial W_2)$ from \mathcal{V}),

respectively. Since h_t is an isotopy defined on \mathcal{V} , considering the latter part of the previous “if and only if” statement, it is equivalent to the statement that V_2 cuts off (i) a solid torus missing ∂W_2 , (ii) a handlebody of genus at least two missing ∂W_2 , or (iii) a compression body with nonempty negative boundary missing ∂W_2 from \mathcal{V} , respectively. This completes the proof of Case b.

This completes the proof of Claim 1. □

Similarly, we get Claim 2.

- CLAIM 2. (i) W_1 is nonseparating or cuts off a solid torus missing ∂V_1 from \mathcal{W} if and only if W_2 is nonseparating or cuts off a solid torus missing ∂V_2 from \mathcal{W} .
(ii) W_1 cuts off a handlebody of genus at least two missing ∂V_1 from \mathcal{W} if and only if W_2 cuts off a handlebody of genus at least two missing ∂V_2 from \mathcal{W} .
(iii) W_1 cuts off a compression body with nonempty negative boundary missing ∂V_1 from \mathcal{W} if and only if W_2 cuts off a compression body with nonempty negative boundary missing ∂V_2 from \mathcal{W} .

Recall that \mathbf{H}_1 and \mathbf{H}_2 are of the same type in the sense of Lemma 3.1 by Lemma 3.3. Hence, considering each case of Definition 3.14 for \mathbf{H}_1 and \mathbf{H}_2 awaring of Claim 1 and Claim 2 (excluding type (d)), we conclude \mathbf{H}_1 and \mathbf{H}_2 are of the same type in the sense of Definition 3.14.

This completes the proof. \square

Considering Lemma 3.15, we can say an equivalent class of \mathcal{GHS}_F^* / \sim is of one type among the ten types in Definition 3.14. Moreover, we can say a \mathcal{V} - or \mathcal{W} -facial cluster having OEC is of one type among these ten types by the second statement of Lemma 3.15 (but we will show that type (a)-(iii), type (b)- \mathcal{W} -(ii), type (b)- \mathcal{V} -(ii), type (c), and type (d) cases do not induce a \mathcal{V} - or \mathcal{W} -facial cluster).

Lemma 3.16. *Let M and F be as in Lemma 3.1. For a k -simplex Σ in $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ having at least one weak reducing pair, if the GHSs obtained by weak reductions along the weak reducing pairs in Σ are all equivalent, then $k \leq 3$. Moreover, if $k = 3$, then $\Sigma = \{V_1, V_2, W_1, W_2\}$, where V_2 and W_2 cut off solid tori \mathcal{V}' and \mathcal{W}' from \mathcal{V} and \mathcal{W} , respectively, and V_1 and W_1 are meridian disks of \mathcal{V}' and \mathcal{W}' , respectively. This means if $k \geq 4$, then there are at least two weak reducing pairs in Σ such that they give nonequivalent GHSs after weak reductions.*

Proof. Suppose the GHSs obtained by weak reductions along the weak reducing pairs in the k -simplex Σ are all equivalent, where $\Sigma = \{V_1, \dots, V_m, W_1, \dots, W_n\}$, where $V_i \subset \mathcal{V}$ for $1 \leq i \leq m$ ($1 \leq m$), $W_i \subset \mathcal{W}$ for $1 \leq i \leq n$ ($1 \leq n$), and $m + n = k + 1$. Assume $m > 2$. Then each of the three \mathcal{V} -faces $\Delta_1 = \{V_1, V_2, W_1\}$, $\Delta_2 = \{V_2, V_3, W_1\}$ and $\Delta_3 = \{V_1, V_3, W_1\}$ has OEC by the assumption. Considering Δ_1 and Δ_2 , V_2 is nonseparating and V_1 and V_3 are separating in \mathcal{V} by Lemma 3.7. But considering Δ_2 and Δ_3 , V_3 is nonseparating in \mathcal{V} by Lemma 3.7, leading to a contradiction. Hence, we get $m \leq 2$. Similarly, we get $n \leq 2$ and therefore $k + 1 = m + n \leq 4$, i.e. $k \leq 3$. The second statement is obtained by applying Lemma 3.5 to the \mathcal{V} -face $\{V_1, V_2, W_1\}$ and the \mathcal{W} -face $\{V_1, W_1, W_2\}$, where each of them has OEC by the assumption. This completes the proof. \square

Lemma 3.17. *Let M and F be as in Lemma 3.1. Let $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ be a \mathcal{V} -facial cluster having OEC and a \mathcal{W} -facial cluster having OEC, respectively, such that they share the common center. Then $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ have the same equivalent class of type (a)-(i).*

Proof. Considering Lemma 3.7, the GHSs obtained by weak reductions along the weak reducing pairs in $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ are all equivalent to the GHS corresponding to the common center. Moreover, they are of the same type in the sense of Definition 3.14 by Lemma 3.15. Since

$\varepsilon_{\mathcal{V}}$ is a \mathcal{V} -facial cluster having OEC and $\varepsilon_{\mathcal{W}}$ is a \mathcal{W} -facial cluster having OEC, the common center consists of nonseparating disks by Definition 3.8, i.e. the center gives a GHS of type (a)-(i) or type (d) after weak reduction by Definition 3.14. But every weak reducing pair in $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ other than the common center has a separating disk by Definition 3.8 and therefore it cannot give a type (d) GHS after weak reduction by Lemma 3.1. Hence, we conclude both $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ are of type (a)-(i). This completes the proof. \square

Lemma 3.18. *Let M and F be as in Lemma 3.1 and let (V, W) be a weak reducing pair such that the GHS obtained by weak reduction along (V, W) is of type (a)-(i). Then there are a naturally determined \mathcal{V} -facial cluster having OEC, say $\varepsilon_{\mathcal{V}}$, and a naturally determined \mathcal{W} -facial cluster having OEC, say $\varepsilon_{\mathcal{W}}$, satisfying the following:*

- (1) $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ share the common center.
- (2) Let Σ be the union of all simplices of $\mathcal{D}(F)$ spanned by the vertices of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$. Then Σ contains (V, W) .
- (3) The GHSs obtained by weak reductions along the weak reducing pairs in Σ are all equivalent.

Proof. Take \bar{V} (resp. \bar{W}) as V (resp. W) itself if V (resp. W) is nonseparating. If V is separating, then it cuts off a solid torus \mathcal{V}' from \mathcal{V} such that $\partial W \cap \mathcal{V}' = \emptyset$ by Definition 3.14 and therefore we can choose a meridian disk \bar{V} of \mathcal{V}' missing $V \cup W$. Similarly, if W is separating, then it cuts off a solid torus \mathcal{W}' from \mathcal{W} such that $\partial V \cap \mathcal{W}' = \emptyset$ and therefore we can choose a meridian disk \bar{W} of \mathcal{W}' missing $V \cup W$. If both V and W are separating, then $\mathcal{V}' \cap \mathcal{W}' = \emptyset$ because the assumption that $\partial W \cap \mathcal{V}' = \emptyset$ and $\partial V \cap \mathcal{W}' = \emptyset$ in Definition 3.14 means the two once-punctured tori cut by ∂W and ∂V from F are pairwise disjoint, i.e. the four disks V, \bar{V}, \bar{W} and W are pairwise disjoint.

Hence, we can find a weak reducing pair (\bar{V}, \bar{W}) such that either

- (1) (V, W) is (\bar{V}, \bar{W}) itself,
- (2) (V, W) is contained in a \mathcal{V} -face $\Delta_{\mathcal{V}} = \{V, \bar{V}, \bar{W} = W\}$ and it has OEC by Lemma 3.5,
- (3) (V, W) is contained in a \mathcal{W} -face $\Delta_{\mathcal{W}} = \{\bar{V} = V, \bar{W}, W\}$ and it has OEC by Lemma 3.5, or
- (4) (V, W) is contained in a 3-simplex $\Sigma_{VW} = \{V, \bar{V}, \bar{W}, W\}$ such that \bar{V} and \bar{W} are meridian disks of the solid tori that V and W cut off from \mathcal{V} and \mathcal{W} , respectively.


Here, (\bar{V}, \bar{W}) is uniquely determined in $\mathcal{D}(F)$ by (V, W) because the meridian disk of a solid torus is unique up to isotopy.

From now on, if we say a 3-simplex Σ_{VW} satisfies the condition $(*)$ in the proof of this lemma, then it will mean that $V \subset \mathcal{V}$ and $W \subset \mathcal{W}$ cut off solid tori from \mathcal{V} and \mathcal{W} , respectively, and the other two disks of Σ_{VW} are meridian disks of the solid tori, respectively, as in the case (4), where it is the same condition for the 3-simplex Σ in the second statement of Lemma 3.16.

CLAIM 3. (V, W) is contained in a 3-simplex of the form

$$\Sigma_{V'W'} = \{V', \bar{V}, \bar{W}, W'\}$$

satisfying the condition $(*)$ in any case as well as the case (4).

Fig.3. $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$

Proof of Claim 3. Let us consider the case (2). Then (i) ∂V divides F into a once-punctured genus at least two surface F' and a once-punctured torus F'' and (ii) $\partial \bar{V} \subset F''$ and $\partial \bar{W} \subset F'$ because $\partial \bar{W} (= \partial W)$ misses the solid torus \mathcal{V}' . Here, $\partial \bar{W}$ is nonseparating in F' by Lemma 2.14. Hence, we can find a band-sum of two parallel copies of \bar{W} in \mathcal{W} , say W' , such that $\partial W' \subset F'$ (if we use the expression “two parallel copies of \bar{W} in \mathcal{W} ”, then we will assume that there is a product neighborhood $\bar{W} \times I$ of \bar{W} in \mathcal{W} such that the disks $\bar{W} \times \{0, 1\}$ are the two parallel copies of \bar{W} and \bar{W} is $\bar{W} \times \{1/2\}$ and see Figure 2 of [4] for the idea of “band-sum”), i.e. W' cuts off a solid torus \mathcal{W}' from \mathcal{W} missing the \mathcal{V} -disks of $\Delta_{\mathcal{V}}$ and \bar{W} is a meridian disk of \mathcal{W}' . Hence, (V, W) belongs to the 3-simplex $\Sigma_{V'W'} = \{V = V', \bar{V}, W = \bar{W}, W'\}$ satisfying the condition (*). The symmetric argument also holds for the case (3).

Let us consider the case (1). Then we can find a band-sum of two parallel copies of V (resp. W) in \mathcal{V} (resp. \mathcal{W}), say V' (resp. W'), such that $V' \cap W' = \emptyset$ by using two disjoint simple arcs in F connecting two parallel copies of V and W , respectively, because $\partial V \cup \partial W$ is nonseparating in F by Lemma 3.1 and therefore we can find such two disjoint arcs in the connected surface F_{VW} whose genus is at least one, where the arcs connect two scars of V and those of W , respectively. Hence, $\Sigma_{V'W'} = \{V', V = \bar{V}, W = \bar{W}, W'\}$ forms a 3-simplex satisfying the condition (*).

This completes the proof of Claim 3. \square

Hence, considering the \mathcal{V} -face $\{V', \bar{V}, \bar{W}\}$ and the \mathcal{W} -face $\{\bar{V}, \bar{W}, W'\}$ contained in the 3-simplex $\Sigma_{V'W'}$, we can see V and W belong to these \mathcal{V} - and \mathcal{W} -faces, respectively, and each face has OEC by Lemma 3.5. Using Lemma 3.9, we can guarantee the existence of the \mathcal{V} -facial cluster having OEC, say $\varepsilon_{\mathcal{V}}$, and the \mathcal{W} -facial cluster having OEC, say $\varepsilon_{\mathcal{W}}$, such that each of them contains (\bar{V}, \bar{W}) and both V and W belong to $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$. Moreover, the GHSS obtained by weak reductions along the weak reducing pairs in $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ are all equivalent because $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ share (\bar{V}, \bar{W}) .

We can observe the following (see Fig.3 and Definition 3.8):

- (a) Only \bar{V} and \bar{W} are nonseparating disks among the vertices of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$.
- (b) Each of \bar{V} and \bar{W} is connected to every other vertex of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ by a 1-simplex in $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$.

Let Σ be the union of all simplices of $\mathcal{D}(F)$ spanned by the vertices of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ as in the statement of this lemma.

CLAIM 4. There is no 1-simplex in $D(F)$ spanned by two separating disks of $\Sigma \cap D_{\mathcal{V}}(F)$ (resp. $\Sigma \cap D_{\mathcal{W}}(F)$).

Proof of Claim 4. For the sake of contradiction, assume there is a 1-simplex σ in $D(F)$ spanned by two separating disks of $\Sigma \cap D_{\mathcal{V}}(F)$. By (a), the vertices of σ are two disks of $\varepsilon_{\mathcal{V}} \cap D_{\mathcal{V}}(F)$ other than \bar{V} , violating Lemma 3.11. This completes the proof of Claim 4. \square

CLAIM 5. Σ is equal to the union of all $\Sigma_{V'W'}$ satisfying the condition (*) and containing (\bar{V}, \bar{W}) .

Proof of Claim 5. Let us consider the union of all $\Sigma_{V'W'}$ satisfying the condition (*) and containing (\bar{V}, \bar{W}) . Considering the condition (*) for each $\Sigma_{V'W'}$ in the union, V' and W' cut off solid tori from \mathcal{V} and \mathcal{W} , respectively, and \bar{V} and \bar{W} are meridian disks of the solid tori, respectively, and therefore each of the \mathcal{V} -face $\Delta'_{\mathcal{V}} = \{V', \bar{V}, \bar{W}\}$ and the \mathcal{W} -face $\Delta'_{\mathcal{W}} = \{\bar{V}, \bar{W}, W'\}$ has OEC by Lemma 3.5. Considering the \mathcal{V} -facial cluster having OEC $\varepsilon'_{\mathcal{V}}$ which contains $\Delta'_{\mathcal{V}}$, where the existence and uniqueness are guaranteed by Lemma 3.9, the center must be (\bar{V}, \bar{W}) by Definition 3.8 because \bar{V} is nonseparating. But Corollary 3.10 forces $\varepsilon'_{\mathcal{V}}$ to be $\varepsilon_{\mathcal{V}}$ because they share the common center, i.e. $\Delta'_{\mathcal{V}} \subset \varepsilon_{\mathcal{V}}$. Likewise, we get $\Delta'_{\mathcal{W}} \subset \varepsilon_{\mathcal{W}}$. This means $\Sigma_{V'W'}$ itself is a 3-simplex spanned by four vertices of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ and therefore the union belongs to Σ . Hence, we will prove Σ belongs to the union.

It is sufficient to show that each simplex in Σ belongs to some $\Sigma_{V'W'}$ in the union. We will consider 1- or more simplices in Σ . (Since every vertex of Σ is contained in a \mathcal{V} -face of $\varepsilon_{\mathcal{V}}$ or a \mathcal{W} -face of $\varepsilon_{\mathcal{W}}$ which is a 2-simplex in Σ , if we can prove it for every 2-simplex in Σ , then all vertices in Σ would belong to the union.)

Case a. $\sigma \subset \Sigma$ is a 1-simplex.

If σ is (\bar{V}, \bar{W}) , then there is nothing to prove. Assume $\sigma \neq (\bar{V}, \bar{W})$. Suppose σ contains a nonseparating disk, i.e. it consists of \bar{V} or \bar{W} and a separating disk by (a). Since the separating disk of σ is also a vertex of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$, σ is the 1-simplex in $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$ described in (b), i.e. there is a \mathcal{V} - or \mathcal{W} -face having OEC containing σ , say Δ , which is contained in $\varepsilon_{\mathcal{V}}$ or $\varepsilon_{\mathcal{W}}$, respectively, and therefore it also contains the common center (\bar{V}, \bar{W}) . If Δ is a \mathcal{V} -face, then one of the \mathcal{V} -disks of Δ cuts off a solid torus from \mathcal{V} and the other is the meridian disk of it by Lemma 3.5 and vice versa. Hence, if we use the proof of Claim 3 corresponding to the case (2) or (3), then we can find a 3-simplex $\Sigma_{V'W'}$ containing Δ such that it satisfies the condition (*) and contains (\bar{V}, \bar{W}) , leading to the result.

Hence, assume σ consists of two separating disks. If σ is not a weak reducing pair, then this violates Claim 4. Hence, σ is a weak reducing pair, say (V', W') , i.e. we can find a \mathcal{V} -face $\Delta_{\mathcal{V}}^{\sigma} = \{V', \bar{V}, \bar{W}\}$ in $\varepsilon_{\mathcal{V}}$ intersecting σ and a \mathcal{W} -face $\Delta_{\mathcal{W}}^{\sigma} = \{\bar{V}, \bar{W}, W'\}$ in $\varepsilon_{\mathcal{W}}$ intersecting σ . This means $\Delta_{\mathcal{V}}^{\sigma}$, $\Delta_{\mathcal{W}}^{\sigma}$ and σ form a 3-simplex $\Sigma_{V'W'} = \{V', \bar{V}, \bar{W}, W'\}$. Since each of $\Delta_{\mathcal{V}}^{\sigma}$ and $\Delta_{\mathcal{W}}^{\sigma}$ has OEC, considering Lemma 3.5, we conclude $\Sigma_{V'W'}$ satisfies the condition (*), leading to the result.

Case b. $\Delta \subset \Sigma$ is a 2-simplex.

If Δ is not a \mathcal{V} - or \mathcal{W} -face, then either $\Delta \subset D_{\mathcal{V}}(F)$ or $\Delta \subset D_{\mathcal{W}}(F)$. Without loss of

generality, assume $\Delta \subset D_{\mathcal{V}}(F)$. Then there is at least one 1-simplex $\sigma \subset \Delta$ spanned by two separating disks of $\Sigma \cap D_{\mathcal{V}}(F)$ by (a), violating Claim 4.

Hence, Δ is a \mathcal{V} - or \mathcal{W} -face.

If Δ belongs to $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$, then we can find a 3-simplex $\Sigma_{V'W'}$ satisfying the condition (*) and containing Δ by the same argument for Δ in Case a. Since Δ contains (\bar{V}, \bar{W}) , this leads to the result.

If Δ does not belong to $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$, then exactly one of \bar{V} and \bar{W} does not belong to Δ . ((i) If Δ contains both \bar{V} and \bar{W} , then Δ is formed by the common center of $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ and the other disk coming from $\varepsilon_{\mathcal{V}}$ or $\varepsilon_{\mathcal{W}}$, i.e. Δ is a \mathcal{V} -face of $\varepsilon_{\mathcal{V}}$ or a \mathcal{W} -face of $\varepsilon_{\mathcal{W}}$, leading to a contradiction, and (ii) if Δ misses both \bar{V} and \bar{W} , then Δ consists of two separating \mathcal{V} -disks and one separating \mathcal{W} -disk by (a) or vice versa. Then there is a 1-simplex $\sigma \subset \Delta$ spanned by two separating disks of $\Sigma \cap D_{\mathcal{V}}(F)$ or $\Sigma \cap D_{\mathcal{W}}(F)$, respectively, violating Claim 4.)

Without loss of generality, assume $\bar{V} \notin \Delta$ and $\bar{W} \in \Delta$. In this case, Δ must be a \mathcal{W} -face otherwise there is a 1-simplex $\sigma \subset \Delta$ spanned by two separating disks of $\Sigma \cap D_{\mathcal{V}}(F)$ by (a), violating Claim 4. Moreover, considering (b), there is a 1-simplex connecting \bar{V} and each vertex of Δ . This means Δ forms a 3-simplex together with \bar{V} , where this 3-simplex contains (\bar{V}, \bar{W}) and consists of two disks from \mathcal{V} and two disks from \mathcal{W} . Here, the two disks of this 3-simplex other than \bar{V} and \bar{W} cut off solid tori from the relevant compression bodies and \bar{V} and \bar{W} are meridian disks of the solid tori, respectively, by Definition 3.8 (because the vertices come from $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$) and therefore this 3-simplex has the form $\Sigma_{V'W'} = \{V', \bar{V}, \bar{W}, W'\}$ satisfying the condition (*), leading to the result.

Case c. $\Sigma' \subset \Sigma$ is a 3-simplex.

If Σ' contains a 2-simplex Δ which is not a \mathcal{V} - or \mathcal{W} -face, then we get a contradiction as in the start of Case b. This means Σ' has the form $\{V_1, V_2, W_1, W_2\}$ for $V_i \subset \mathcal{V}$ and $W_i \subset \mathcal{W}$ for $i = 1, 2$. Moreover, applying Claim 4 to the 1-simplices $\{V_1, V_2\}$ and $\{W_1, W_2\}$, we can assume $V_1 = \bar{V}$ and $W_1 = \bar{W}$ by (a). Hence, considering the last sentence of Case b, we can see Σ' itself has the form $\Sigma_{V_2W_2} = \{V_2, \bar{V} = V_1, \bar{W} = W_1, W_2\}$ satisfying the condition (*), leading to the result.

If there is a simplex of dimension at least four in Σ , i.e. it consists of at least five disks, then there must be a 2-simplex in Σ which is neither a \mathcal{V} -face nor a \mathcal{W} -face, leading to a contradiction as in the start of Case b. Therefore, we do not need to consider more higher dimensional simplices.

This completes the proof of Claim 5. □

Considering the proof of Claim 5, the 1-simplex (V, W) (which is spanned by two vertices of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$) belongs to Σ . Since the common center (\bar{V}, \bar{W}) of $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ is uniquely determined by (V, W) , $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ are also uniquely determined by (V, W) by Corollary 3.10. This completes the proof of all statements of this Lemma other than (3.18).

CLAIM 6. The GHSs obtained by weak reductions along the weak reducing pairs in a 3-simplex $\Sigma_{V'W'}$ satisfying the condition (*) are all equivalent.

Proof of Claim 6. Let $\Sigma_{V'W'} = \{V', \tilde{V}, \tilde{W}, W'\}$ and suppose (D, E) is a weak reducing pair in $\Sigma_{V'W'}$. We will prove the GHS obtained by weak reduction along (D, E) is equivalent to that obtained by weak reduction along (\tilde{V}, \tilde{W}) .

First, considering $\Sigma_{V'W'}$ satisfies the condition $(*)$, we can see each of the \mathcal{V} -face $\Delta_{\mathcal{V}} = \{V', \tilde{V}, \tilde{W}\}$ and the \mathcal{W} -face $\Delta_{\mathcal{W}} = \{\tilde{V}, \tilde{W}, W'\}$ has OEC by Lemma 3.5.

If one of D and E is \tilde{V} or \tilde{W} , then (D, E) belongs to $\Delta_{\mathcal{W}}$ or $\Delta_{\mathcal{V}}$, respectively, leading to the result.

Suppose $D = V'$ and $E = W'$. In this case, we get the \mathcal{V} -face $\Delta'_{\mathcal{V}} = \{V', \tilde{V}, W'\}$ containing (V', W') and $\Delta'_{\mathcal{V}}$ has OEC by Lemma 3.5. Moreover, $\Delta'_{\mathcal{V}}$ shares the weak reducing pair (\tilde{V}, W') with $\Delta_{\mathcal{W}}$, i.e. (D, E) gives a GHS equivalent to that obtained by using (\tilde{V}, \tilde{W}) after weak reduction.

This completes the proof of Claim 6. \square

Considering the proof of Claim 5, any weak reducing pair (D, E) in Σ belongs to a 3-simplex $\Sigma_{V'W'}$ satisfying the condition $(*)$ and containing (\tilde{V}, \tilde{W}) . Moreover, considering Claim 6, we conclude the GHS obtained by weak reduction along (D, E) is equivalent to the GHS obtained by weak reduction along (\tilde{V}, \tilde{W}) , i.e. the GHSs obtained by weak reductions along the weak reducing pairs in Σ are all equivalent to that corresponding to (\tilde{V}, \tilde{W}) . This completes the proof of (3.18).

This completes the proof of Lemma 3.18. \square

DEFINITION 3.19. Let M and F be as in Lemma 3.1. An *equivalent cluster of type (a)-(i)* is the union of all simplices of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ spanned by the vertices of $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$, where $\varepsilon_{\mathcal{V}}$ is a \mathcal{V} -facial cluster having OEC, $\varepsilon_{\mathcal{W}}$ is a \mathcal{W} -facial cluster having OEC, and they share the common center. We call the common center of $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$ the *center* of the equivalent cluster. Lemma 3.18 means every weak reducing pair giving a type (a)-(i) GHS after weak reduction belongs to a naturally determined equivalent cluster of type (a)-(i).

Lemma 3.20. *An equivalent cluster of type (a)-(i) is contractible.*

Proof. Considering Claim 5, the shape of an equivalent cluster of type (a)-(i) is essentially the same as that of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ of Case (a) in the proof of Lemma 3.5 of [5] and it is contractible. Therefore, we omit the proof. \square

Lemma 3.21. *Let M and F be as in Lemma 3.1 and let (V, W) be a weak reducing pair such that the GHS obtained by weak reduction along (V, W) is of type (a)-(ii)- \mathcal{W} or type (b)- \mathcal{W} -(i). Then there is a naturally determined \mathcal{W} -facial cluster having OEC containing (V, W) . Likewise, if the GHS obtained by weak reduction along (V, W) is of type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i), then there is a naturally determined \mathcal{V} -facial cluster having OEC containing (V, W) .*

Proof. Suppose the GHS obtained by weak reduction along (V, W) is of type (a)-(ii)- \mathcal{W} or type (b)- \mathcal{W} -(i). We can see the following from Definition 3.14:

- (1) W is nonseparating or W cuts off a solid torus \mathcal{W}' missing ∂V from \mathcal{W} .
- (2) Either V cuts off a handlebody of genus at least two from \mathcal{V} missing ∂W (the type (a)-(ii)- \mathcal{W} case) or a compression body with nonempty negative boundary missing

∂W from \mathcal{V} (the type (b)- \mathcal{W} -(i) case).

Let \bar{V} be V itself. Take \bar{W} as W itself if W is nonseparating in \mathcal{W} . If W is separating, then we take \bar{W} as a meridian disk of \mathcal{W}' . Hence, we can find a weak reducing pair (\bar{V}, \bar{W}) such that either

- (1) (V, W) is (\bar{V}, \bar{W}) itself, or
- (2) (V, W) is contained in the \mathcal{W} -face $\Delta_{\mathcal{W}} = \{\bar{V}, \bar{W}, W\}$ and $\Delta_{\mathcal{W}}$ has OEC by Lemma 3.5.

Here, (\bar{V}, \bar{W}) is uniquely determined in $\mathcal{D}(F)$ by (V, W) because the meridian disk of a solid torus is unique up to isotopy. Hence, it is sufficient to show that there exists a \mathcal{W} -facial cluster having OEC determined by (V, W) such that it contains (V, W) and its center is (\bar{V}, \bar{W}) by Corollary 3.10. By Lemma 3.9, the proof for the case (2) is obvious.

Consider the case (1). Then $\partial \bar{W}$ is nonseparating in the relevant component of $F_{\bar{V}}$ by Lemma 2.14, where this component is of genus at least two by Lemma 2.3, and therefore we can take a band-sum of two parallel copies of $W = \bar{W}$ in \mathcal{W} , say W' , such that $\Delta = \{V = \bar{V}, W = \bar{W}, W'\}$ is a \mathcal{W} -face. Moreover, Δ has OEC by Lemma 3.5. Hence, we can guarantee the existence of a \mathcal{W} -facial cluster having OEC and containing (V, W) by Lemma 3.9. Since W is nonseparating, (\bar{V}, \bar{W}) is the center of the \mathcal{W} -facial cluster.

This completes the proof. \square

DEFINITION 3.22. Let M and F be as in Lemma 3.1. Let $\varepsilon_{\mathcal{W}}$ be a \mathcal{W} -facial cluster having OEC such that the GHSs obtained by weak reductions along the weak reducing pairs in $\varepsilon_{\mathcal{W}}$ are either of type (a)-(ii)- \mathcal{W} or type (b)- \mathcal{W} -(i). We call $\varepsilon_{\mathcal{W}}$ an *equivalent cluster of type (a)-(ii)- \mathcal{W} or (b)- \mathcal{W} -(i)*, respectively. We call the center of $\varepsilon_{\mathcal{W}}$ the *center* of the equivalent cluster.

Lemma 3.21 means every weak reducing pair giving a type (a)-(ii)- \mathcal{W} or (b)- \mathcal{W} -(i) GHS after weak reduction belongs to a naturally determined equivalent cluster of type (a)-(ii)- \mathcal{W} or type (b)- \mathcal{W} -(i), respectively.

Likewise, we get the following definition.

DEFINITION 3.23. Let M and F be as in Lemma 3.1. Let $\varepsilon_{\mathcal{V}}$ be a \mathcal{V} -facial cluster having OEC such that the GHSs obtained by weak reductions along the weak reducing pairs in $\varepsilon_{\mathcal{V}}$ are either of type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i). We call $\varepsilon_{\mathcal{V}}$ an *equivalent cluster of type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i)*, respectively. We call the center of $\varepsilon_{\mathcal{V}}$ the *center* of the equivalent cluster.

Lemma 3.21 means every weak reducing pair giving a type (a)-(ii)- \mathcal{V} or (b)- \mathcal{V} -(i) GHS after weak reduction belongs to a naturally determined equivalent cluster of type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i), respectively.

Since a \mathcal{W} - or \mathcal{V} -facial cluster having OEC is contractible, an equivalent cluster of type (a)-(ii)- \mathcal{W} , type (b)- \mathcal{W} -(i), type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i) is contractible. Moreover, Lemma 3.13 means an equivalent cluster of type (a)-(ii)- \mathcal{W} , type (b)- \mathcal{W} -(i), type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i) is the union of all simplices of $\mathcal{D}(F)$ spanned by the vertices of the cluster.

DEFINITION 3.24. Let M and F be as in Lemma 3.1. Suppose the GHS obtained by weak reduction along (V, W) is of type (a)-(iii), type (b)- \mathcal{W} -(ii), type (b)- \mathcal{V} -(ii), type (c), or type




Fig.4. the shapes of equivalent clusters

(d). In this case, we call the weak reducing pair (V, W) itself *an equivalent cluster of type (a)-(iii), type (b)- \mathcal{W} -(ii), type (b)- \mathcal{V} -(ii), type (c), or type (d)*, respectively. We also define the *center* of the equivalent cluster (\bar{V}, \bar{W}) as (V, W) itself.

See Fig.4 for the shapes of equivalent clusters.

In summary, we get the following two lemmas.

First, we can understand the *type* of an equivalent cluster as the type of GHSs obtained by weak reductions in the cluster and therefore we summarize all cases in Lemma 3.25. Since any weak reducing pair gives exactly one type of GHS after weak reduction in the sense of Definition 3.14, the second statement of Lemma 3.25 comes from Lemma 3.15 directly.

Lemma 3.25. *Let M and F be as in Lemma 3.1 and let \mathcal{B} be an equivalent cluster. Then the GHSs obtained by weak reductions along the weak reducing pairs in \mathcal{B} are all equivalent. Moreover, the GHSs are of the same type in the sense of Definition 3.14, where the type is uniquely determined.*

Next, Lemma 3.26 means every equivalent cluster is contractible and the intersection of an equivalent cluster and $\mathcal{D}_{\mathcal{V}}(F)$ or $\mathcal{D}_{\mathcal{W}}(F)$ is contractible.

Lemma 3.26. *Let M and F be as in Lemma 3.1. Then every equivalent cluster is contractible. Moreover, every equivalent cluster is the union of all simplices of $\mathcal{D}(F)$ spanned by the vertices of the cluster. When an equivalent cluster intersects $\mathcal{D}_{\mathcal{V}}(F)$ or $\mathcal{D}_{\mathcal{W}}(F)$, the dimension of the intersection is at most one. Indeed, the intersection is either a vertex coming from the center if the dimension is zero or a star-shaped graph with infinitely many edges if the dimension is one. Moreover, if the dimension of the intersection is one, then the star-shaped graph is equal to the intersection of the \mathcal{V} - or \mathcal{W} -facial cluster having OEC in the equivalent cluster whose center is the center of the equivalent cluster and $\mathcal{D}_{\mathcal{V}}(F)$ or $\mathcal{D}_{\mathcal{W}}(F)$,*

respectively, i.e. the vertex positioned at the center of the star-shaped graph comes from the center of the equivalent cluster.

Considering Lemma 3.18, Lemma 3.21 and Definition 3.24, any weak reducing pair (V, W) belongs to a naturally determined equivalent cluster, say the *canonical equivalent cluster* for (V, W) .

The next lemma gives a criterion to determine whether a disk in an equivalent cluster belongs to the center of the cluster or not.

Lemma 3.27. *Let M and F be as in Lemma 3.1 and D is a disk in an equivalent cluster \mathcal{B} . Then D belongs to the center of \mathcal{B} if and only if one of the following holds.*

- (1) *it is nonseparating or*
- (2) *if it is separating, then it does not cut off a solid torus from the relevant compression body.*

Proof. Let (\bar{V}, \bar{W}) be the center of \mathcal{B} . Then the following hold:

- (1) **Type (a)-(i):** \bar{V} and \bar{W} are nonseparating, and another disk cuts off a solid torus from the relevant compression body (see Definition 3.19 and Definition 3.8).
- (2) **Type (a)-(ii)- \mathcal{W} and Type (b)- \mathcal{W} -(i):** \bar{W} is nonseparating and another \mathcal{W} -disk cuts off a solid torus from \mathcal{W} (see Definition 3.22 and Definition 3.8). \bar{V} is separating in \mathcal{V} and it does not cut off a solid torus from \mathcal{V} (see Definition 3.14).
- (3) **Type (a)-(ii)- \mathcal{V} and Type (b)- \mathcal{W} -(i):** Use the symmetric argument of the previous case (see Definition 3.23, Definition 3.8 and Definition 3.14).
- (4) **Type (a)-(iii), Type (b)- \mathcal{W} -(ii), Type (b)- \mathcal{V} -(ii) and Type (c):** Each of \bar{V} and \bar{W} is separating, but it does not cut off a solid torus from the relevant compression body (see Lemma 3.1, Definition 3.14 and Definition 3.24).
- (5) **Type (d):** \bar{V} and \bar{W} are nonseparating (see Lemma 3.1 and Definition 3.24).

This completes the proof. \square

Lemma 3.28. *Let M and F be as in Lemma 3.1. Let \mathcal{B}_i be an equivalent cluster and (\bar{V}_i, \bar{W}_i) the center of \mathcal{B}_i for $i = 1, 2$. If $(\mathcal{B}_1 \cap \mathcal{B}_2) \cap \mathcal{D}_{\mathcal{V}}(F) \neq \emptyset$, then $\bar{V}_1 = \bar{V}_2$. Likewise, if $(\mathcal{B}_1 \cap \mathcal{B}_2) \cap \mathcal{D}_{\mathcal{W}}(F) \neq \emptyset$, then $\bar{W}_1 = \bar{W}_2$.*

Proof. Let V be a vertex of $(\mathcal{B}_1 \cap \mathcal{B}_2) \cap \mathcal{D}_{\mathcal{V}}(F)$.

Assume $V = \bar{V}_1$. Then (1) V is nonseparating or (2) it does not cut off a solid torus from \mathcal{V} if it is separating as a disk in \mathcal{B}_1 by Lemma 3.27. Since V also belongs to \mathcal{B}_2 , $V = \bar{V}_2$ by Lemma 3.27, i.e. $\bar{V}_1 = \bar{V}_2$.

Assume $V \neq \bar{V}_1$. Then V cuts off a solid torus \mathcal{V}' from \mathcal{V} as a disk in \mathcal{B}_1 by Lemma 3.27 and therefore $V \neq \bar{V}_2$ as a disk in \mathcal{B}_2 by Lemma 3.27. Here, each \mathcal{B}_i contains a \mathcal{V} -facial cluster having OEC whose center is (\bar{V}_i, \bar{W}_i) , say $\mathcal{E}_{\mathcal{V}}^i$, and V is a vertex of the star-shaped graph $\mathcal{E}_{\mathcal{V}}^i \cap \mathcal{D}_{\mathcal{V}}(F)$ which is not the center of the graph by Lemma 3.26 for $i = 1, 2$. This means \bar{V}_1 and \bar{V}_2 are meridian disks of the solid torus \mathcal{V}' by Definition 3.8, i.e. they are isotopic in \mathcal{V} . Therefore, we get $\bar{V}_1 = \bar{V}_2$.

This completes the proof. \square

The next lemma means the canonical equivalent cluster for (V, W) is the only equivalent cluster containing (V, W) .

Lemma 3.29. *Let M and F be as in Lemma 3.1. Then every weak reducing pair belongs to a uniquely determined equivalent cluster. This means if two different equivalent clusters intersect each other, then the intersection cannot contain a weak reducing pair.*

Proof. By definitions of equivalent clusters, every weak reducing pair belongs to some equivalent cluster. Let us consider a weak reducing pair (V, W) . Suppose there are two equivalent clusters containing (V, W) , say B_1 and B_2 . Let (\bar{V}_i, \bar{W}_i) be the center of B_i for $i = 1, 2$. We will prove $B_1 = B_2$.

CLAIM 7. B_1 and B_2 are of the same type and they share the common center.

Proof of Claim 7. Since B_1 and B_2 share (V, W) , the GHSs obtained by weak reductions along the weak reducing pairs in $B_1 \cup B_2$ are all equivalent to that corresponding to (V, W) and they are of the same type in the sense of Definition 3.14 by Lemma 3.25. This means B_1 and B_2 are of the same type.

Since B_1 and B_2 share (V, W) , $(B_1 \cap B_2) \cap D_V(F) \neq \emptyset$ and $(B_1 \cap B_2) \cap D_W(F) \neq \emptyset$. Therefore, Lemma 3.28 means $\bar{V}_1 = \bar{V}_2$ and $\bar{W}_1 = \bar{W}_2$, i.e. $(\bar{V}_1, \bar{W}_1) = (\bar{V}_2, \bar{W}_2)$.

This completes the proof of Claim 7. \square

By Claim 7, B_1 and B_2 are of the same type (so the shapes of them are the same in the sense of Fig.4) and they share the common center. Hence, if each of B_1 and B_2 consists of a weak reducing pair, then we conclude $B_1 = B_2$, leading to the result. Hence, assume each of B_1 and B_2 does not consist of a weak reducing pair. Considering $B_i \cap D_V(F)$ and $B_i \cap D_W(F)$, if both are of dimension zero, then each of them consists of a vertex by Lemma 3.26, i.e. B_i is a weak reducing pair, leading to a contradiction. Hence, at least one of $B_i \cap D_V(F)$ and $B_i \cap D_W(F)$ is of dimension one by Lemma 3.26 for $i = 1, 2$.

If $\dim(B_i \cap D_V(F)) = 1$ for $i = 1, 2$, then each of B_1 and B_2 contains a \mathcal{V} -facial cluster having OEC whose center is (\bar{V}_i, \bar{W}_i) , say ε_V^i , by Lemma 3.26 for $i = 1, 2$. Here, Claim 7 means the center of ε_V^1 is equal to that of ε_V^2 and therefore we get $\varepsilon_V^1 = \varepsilon_V^2$ by Corollary 3.10. Hence, say $\varepsilon_V := \varepsilon_V^1 = \varepsilon_V^2$ in this case.

Likewise, if $\dim(B_i \cap D_W(F)) = 1$ for $i = 1, 2$, then each of B_1 and B_2 contains a \mathcal{W} -facial cluster having OEC whose center is (\bar{V}_i, \bar{W}_i) , say ε_W^i , and therefore we get $\varepsilon_W^1 = \varepsilon_W^2$ likewise. Hence, say $\varepsilon_W := \varepsilon_W^1 = \varepsilon_W^2$ in this case.

Considering the uniqueness of the \mathcal{V} - or \mathcal{W} -facial cluster having OEC whose center is (\bar{V}_i, \bar{W}_i) by Corollary 3.10, we can see either

- (1) $\varepsilon_V \cup \varepsilon_W$ determines B_i as in Definition 3.19 (B_i is of type (a)-(i)) for $i=1,2$,
- (2) ε_W is equal to B_i as in Definition 3.22 (B_i is of type (a)-(ii)- \mathcal{W} or type (b)- \mathcal{W} -(i)) for $i=1,2$, or
- (3) ε_V is equal to B_i as in Definition 3.23 (B_i is of type (a)-(ii)- \mathcal{V} or type (b)- \mathcal{V} -(i)) for $i=1,2$.

Hence, we conclude $B_1 = B_2$.

This completes the proof. \square

DEFINITION 3.30. Considering Lemma 3.25 and Lemma 3.29, there is the *canonical function* Φ from the set of equivalent clusters to the set of equivalence classes \mathcal{GHS}_F^* / \sim , where

$\Phi(\mathcal{B})$ is the equivalent class of the GHSs obtained by weak reductions along the weak reducing pairs in \mathcal{B} for a given equivalent cluster \mathcal{B} . For a representative $\mathbf{H} \in \mathcal{GHS}_F^*$ of an equivalent class $[\mathbf{H}]^*$, there exists a weak reducing pair giving \mathbf{H} after weak reduction. Moreover, this weak reducing pair belongs to an equivalent cluster \mathcal{B}' by Lemma 3.29, i.e. $\Phi(\mathcal{B}') = [\mathbf{H}]^*$, and therefore Φ is surjective.

As well as a weak reducing pair, a \mathcal{V} - or \mathcal{W} -facial cluster having OEC also belongs to a uniquely determined equivalent cluster by Lemma 3.31.

Lemma 3.31. *Let M and F be as in Lemma 3.1. If there is a \mathcal{V} - or \mathcal{W} -facial cluster having OEC, then it belongs to a uniquely determined equivalent cluster. Therefore, every \mathcal{V} - or \mathcal{W} -face having OEC belongs to a uniquely determined equivalent cluster.*

Proof. Without loss of generality, assume ε is a \mathcal{V} -facial cluster having OEC. Let (V, W) be the center of ε . Then V is nonseparating in \mathcal{V} by Definition 3.8.

CLAIM 8. (V, W) itself is not an equivalent cluster.

Proof of Claim 8. For the sake of contradiction, suppose (V, W) is an equivalent cluster. Let us consider the cases where a weak reducing pair itself is an equivalent cluster. In the cases of type (a)-(iii), type (b)- \mathcal{W} -(ii), type (b)- \mathcal{V} -(ii) and type (c), both disks are separating by Definition 3.14. Therefore, considering that V is nonseparating in \mathcal{V} , (V, W) is of type (d), i.e. both V and W are nonseparating in \mathcal{V} and \mathcal{W} , respectively, and $\partial V \cup \partial W$ is separating in F .

Let D be another disk of ε other than V and W . Then D cuts off a solid torus \mathcal{V}' from \mathcal{V} and V is a meridian disk of \mathcal{V}' by Definition 3.8, i.e. ∂V is nonseparating in the relevant component of F_D by Lemma 2.14. Moreover, ∂W misses \mathcal{V}' by Lemma 2.3 and ∂W is nonseparating in the relevant component of F_D by Lemma 2.14. Let us compress F_D along V and W . Then the resulting one $(F_D)_{VW}$ consists of two components, i.e. F_{VW} consists of only one component by recovering F_{VW} from the two components of $(F_D)_{VW}$. This means $\partial V \cup \partial W$ is nonseparating in F , violating the assumption.

This completes the proof of Claim 8. □

By Claim 8, (V, W) itself cannot be an equivalent cluster, i.e. the GHS obtained by weak reduction along (V, W) is either of type (a)-(i), type (a)-(ii)- \mathcal{W} , type (a)-(ii)- \mathcal{V} , type (b)- \mathcal{W} -(i) or type (b)- \mathcal{V} -(i). Moreover, we can exclude the cases of type (a)-(ii)- \mathcal{W} and type (b)- \mathcal{W} -(i) because V is nonseparating in \mathcal{V} . Hence, considering the proofs of Lemma 3.18 or Lemma 3.21, there is a weak reducing pair (\bar{V}, \bar{W}) uniquely determined by (V, W) such that it becomes the center of an equivalent cluster \mathcal{B} of type (a)-(i) (if W is nonseparating or cuts off a solid torus from \mathcal{W}), type (a)-(ii)- \mathcal{V} (if W cuts off a handlebody of genus at least two missing ∂V from \mathcal{W}) or type (b)- \mathcal{V} -(i) (if W cuts off a compression body with nonempty negative boundary missing ∂V from \mathcal{W}) and \mathcal{B} contains (V, W) .

Since there cannot be two different equivalent clusters sharing (V, W) by Lemma 3.29, if there is an equivalent cluster containing ε , then it must be the only one. Therefore, we only need to prove ε belongs to \mathcal{B} .

Since V is nonseparating in \mathcal{V} and $V \in \mathcal{B}$, V belongs the center of \mathcal{B} by Lemma 3.27.

Suppose W does not cut off a solid torus from \mathcal{W} , i.e. (V, W) is the center of \mathcal{B} as well as

that of ε by Lemma 3.27. In this case, the \mathcal{V} -facial cluster $\varepsilon_{\mathcal{V}}$ forming \mathcal{B} in Definition 3.19 or Definition 3.23 must be equal to ε by Corollary 3.10, i.e. $\varepsilon \subset \mathcal{B}$, leading to the result.

Suppose W cuts off a solid torus \mathcal{W}' from \mathcal{W} , i.e. the case where \mathcal{B} is of a type (a)-(i). Let \tilde{W} be a meridian disk of \mathcal{W}' missing W . Then $V = \bar{V}$ and $\tilde{W} = \bar{W}$ by the proof of Lemma 3.18 and \tilde{W} misses all vertices of ε because every \mathcal{V} -disks of ε cannot intersect the once-punctured torus component of $F - \partial W$ by Lemma 2.3. This means each \mathcal{V} -face $\{V', V, W\}$ of ε belongs to the 3-simplex of the form $\Sigma_{V'W'} = \{V', \bar{V} = V, \bar{W} = \tilde{W}, W' = W\}$, where V' cuts off a solid torus from \mathcal{V} and V is a meridian disk of the solid torus by Definition 3.8. Since \mathcal{B} contains all such 3-simplices by Claim 5, we conclude ε belongs to \mathcal{B} , leading to the result.

Since a \mathcal{V} -face having OEC belongs to a uniquely determined \mathcal{V} -facial cluster having OEC by Lemma 3.9, the second statement comes from the first statement directly.

This completes the proof of Lemma 3.31. \square

Lemma 3.32. *Let M and F be as in Lemma 3.1. For a 3-simplex Σ of $\mathcal{D}(F)$ containing at least one weak reducing pair, if the GHSs obtained by weak reductions along the weak reducing pairs in Σ are all equivalent, then Σ belongs to a uniquely determined equivalent cluster of type (a)-(i).*

Proof. Considering Lemma 3.16, Σ has the form $\{V', V, W, W'\}$, where V' and W' cut off solid tori \mathcal{V}' and \mathcal{W}' from \mathcal{V} and \mathcal{W} , respectively, and V and W are meridian disks of \mathcal{V}' and \mathcal{W}' , respectively, and therefore each of the \mathcal{V} -face $\Delta_{\mathcal{V}} = \{V', V, W\}$ and the \mathcal{W} -face $\Delta_{\mathcal{W}} = \{V, W, W'\}$ has OEC by Lemma 3.5. This gives the \mathcal{V} - and \mathcal{W} -facial clusters having OEC containing $\Delta_{\mathcal{V}}$ and $\Delta_{\mathcal{W}}$, say $\varepsilon_{\mathcal{V}}$ and $\varepsilon_{\mathcal{W}}$, respectively, by Lemma 3.9, and they share the common center (V, W) by Definition 3.8. Using $\varepsilon_{\mathcal{V}} \cup \varepsilon_{\mathcal{W}}$, we can find an equivalent cluster \mathcal{B} of type (a)-(i) by Definition 3.19 and \mathcal{B} contains Σ by Claim 5.

If there is another equivalent cluster containing Σ , say \mathcal{B}' , then \mathcal{B} and \mathcal{B}' share the weak reducing pairs in Σ , violating Lemma 3.29.

This completes the proof. \square

DEFINITION 3.33. Let M and F be as in Lemma 3.1. Let Σ be a simplex of $\mathcal{D}(F)$ containing at least one weak reducing pair. If there is only OEC of the GHSs obtained by weak reductions along the weak reducing pairs in Σ , then we call Σ an *equivalent simplex*. If there are two or more equivalent classes of the GHSs obtained by weak reductions along the weak reducing pairs in Σ , then we call Σ a *nonequivalent simplex*.

In summary, we get the following theorem.

Theorem 3.34. *Let M and F be as in Lemma 3.1. Then $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ consists of equivalent clusters and nonequivalent simplices. Moreover, each equivalent simplex belongs to a uniquely determined equivalent cluster.*

Proof. Recall that $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ consists of all simplices of $\mathcal{D}(F)$ intersecting both $\mathcal{D}_{\mathcal{V}}(F)$ and $\mathcal{D}_{\mathcal{W}}(F)$, i.e. it is the union of all equivalent simplices and all nonequivalent simplices.

Considering Lemma 3.29, Lemma 3.31 and Lemma 3.32, every equivalent simplex of dimension at most three belongs to a uniquely determined equivalent cluster. Moreover, Lemma 3.16 means there is no equivalent simplex of dimension at least four. Hence, all

equivalent simplices are contained in the union of all equivalent clusters. Therefore, the union of all equivalent clusters and all nonequivalent simplices covers $D_{\mathcal{V}\mathcal{W}}(F)$. Since every equivalent cluster is a subset of $D_{\mathcal{V}\mathcal{W}}(F)$, we conclude the previous union is exactly the same as $D_{\mathcal{V}\mathcal{W}}(F)$.

This completes the proof. \square

4. The proof of Theorem 1.2

In this section, we will prove Theorem 1.2.

The next lemma describes the way how two different equivalent clusters intersect each other.

Lemma 4.1. *Suppose M is an irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ is a weakly reducible, unstabilized Heegaard splitting of M of genus $n \geq 3$. If two different equivalent clusters B_1 and B_2 intersect each other, then the center of B_1 intersects that of B_2 in a vertex.*

Proof. Let (\bar{V}_i, \bar{W}_i) be the center of B_i for $i = 1, 2$. If $(B_1 \cap B_2) \cap D_{\mathcal{V}}(F) \neq \emptyset$, then $\bar{V}_1 = \bar{V}_2$ by Lemma 3.28. Likewise, if $(B_1 \cap B_2) \cap D_{\mathcal{W}}(F) \neq \emptyset$, then $\bar{W}_1 = \bar{W}_2$. But at least one of $(B_1 \cap B_2) \cap D_{\mathcal{V}}(F)$ and $(B_1 \cap B_2) \cap D_{\mathcal{W}}(F)$ must be empty otherwise $(\bar{V}_1, \bar{W}_1) = (\bar{V}_2, \bar{W}_2)$, violating Lemma 3.29.

This completes the proof. \square

Lemma 4.2. *Let M and F be as in Lemma 4.1. Then every component of $D_{\mathcal{V}\mathcal{W}}(F)$ is an equivalent cluster if and only if every \mathcal{V} - or \mathcal{W} -face has OEC.*

Proof. (\Rightarrow) Suppose every component of $D_{\mathcal{V}\mathcal{W}}(F)$ is an equivalent cluster. Let Δ be a \mathcal{V} -face $\{V, V', W\}$ without loss of generality. Then Δ belong to an equivalent cluster B by the assumption, i.e. $\{V, V'\} = \Delta \cap D_{\mathcal{V}}(F) \subset B \cap D_{\mathcal{V}}(F)$. Considering Lemma 3.26, there is a \mathcal{V} -facial cluster having OEC in B , say $\varepsilon_{\mathcal{V}}$, such that $\varepsilon_{\mathcal{V}} \cap D_{\mathcal{V}}(F) = B \cap D_{\mathcal{V}}(F)$ and the intersection is a star-shaped graph. This means the 1-simplex $\{V, V'\}$ is an edge of the star-shaped graph $\varepsilon_{\mathcal{V}} \cap D_{\mathcal{V}}(F)$, i.e. $\{V, V'\}$ belongs to a \mathcal{V} -face having OEC in $\varepsilon_{\mathcal{V}}$, say $\Delta_{\mathcal{V}}$. Therefore, we can assume V' cuts off a solid torus \mathcal{V}' from \mathcal{V} and V is a meridian disk of \mathcal{V}' by Lemma 3.5. Hence, Δ also has OEC by Lemma 3.5.

(\Leftarrow) Suppose every \mathcal{V} - or \mathcal{W} -face has OEC.

CLAIM 9. There is no simplex of dimension at least four in $D_{\mathcal{V}\mathcal{W}}(F)$. Moreover, if there is a 3-simplex in $D_{\mathcal{V}\mathcal{W}}(F)$, then it has the form in the statement of Lemma 3.16.

Proof. Note that every highest dimensional simplex of $D_{\mathcal{V}\mathcal{W}}(F)$ must intersect both $D_{\mathcal{V}}(F)$ and $D_{\mathcal{W}}(F)$.

Suppose $\dim(D_{\mathcal{V}\mathcal{W}}(F)) = k$ and there is a k -simplex $\Sigma = \{V_1, \dots, V_m, W_1, \dots, W_n\} \subset D_{\mathcal{V}\mathcal{W}}(F)$, where $V_i \subset \mathcal{V}$ for $1 \leq i \leq m$ ($1 \leq m$), $W_i \subset \mathcal{W}$ for $1 \leq i \leq n$ ($1 \leq n$), and $m + n = k + 1$. Considering the assumption that every \mathcal{V} - or \mathcal{W} -face has OEC and the arguments in the proof of Lemma 3.16, we get $m, n \leq 2$ and confirm the desired shape of a 3-simplex in $D_{\mathcal{V}\mathcal{W}}(F)$.

This completes the proof of Claim 9. \square

CLAIM 10. Every simplex of $D(F)$ intersecting both $D_{\mathcal{V}}(F)$ and $D_{\mathcal{W}}(F)$ is contained in an equivalent cluster.

Proof of Claim 10. By Claim 9, it is sufficient to consider simplices of dimension at most three. Considering Lemma 3.29, every weak reducing pair belongs to an equivalent cluster. Moreover, the assumption that every \mathcal{V} - or \mathcal{W} -face has OEC induces that every \mathcal{V} - or \mathcal{W} -face belongs to an equivalent cluster by Lemma 3.31. Suppose Σ is a 3-simplex intersecting both $D_{\mathcal{V}}(F)$ and $D_{\mathcal{W}}(F)$. By Claim 9, Σ has the form in the statement of Lemma 3.16. Hence, considering the argument in the proof of Lemma 3.32, we can find an equivalent cluster containing Σ .

This completes the proof of Claim 10. \square

Let C be a component of $D_{\mathcal{V}\mathcal{W}}(F)$. Then $C = \cup_{\alpha \in \mathcal{A}} \sigma_{\alpha}$, where each σ_{α} is a simplex intersecting both $D_{\mathcal{V}}(F)$ and $D_{\mathcal{W}}(F)$ by the definition of $D_{\mathcal{V}\mathcal{W}}(F)$ and \mathcal{A} is an index set. Hence, considering Claim 10, $C \subset \cup_{\alpha \in \mathcal{A}} B_{\alpha}$, where B_{α} is an equivalent cluster containing σ_{α} . Moreover, every equivalent cluster is a connected subset of $D_{\mathcal{V}\mathcal{W}}(F)$ and therefore $B_{\alpha} \subset C$ for all $\alpha \in \mathcal{A}$, i.e. $\cup_{\alpha \in \mathcal{A}} B_{\alpha} \subset C$, and therefore we conclude $C = \cup_{\alpha \in \mathcal{A}} B_{\alpha}$.

It is sufficient to show that C consists of only one B_{α} . For the sake of contradiction, assume there exist B_{α_1} and B_{α_2} such that $B_{\alpha_1} \neq B_{\alpha_2}$. If $B_{\alpha} \cap B_{\beta} = \emptyset$ for all mutually different B_{α} and B_{β} , then C is disconnected, leading to a contradiction, i.e. we can assume $B_{\alpha_1} \cap B_{\alpha_2} \neq \emptyset$. Let (\bar{V}_i, \bar{W}_i) be the center of B_{α_i} for $i = 1, 2$. Then (\bar{V}_1, \bar{W}_1) intersects (\bar{V}_2, \bar{W}_2) in a vertex by Lemma 4.1. Therefore, there is a sequence of \mathcal{V} - and \mathcal{W} -faces $\Delta_0, \Delta_1, \dots, \Delta_n$ such that the following hold by Lemma 8.4 of [1]:

- (1) $\Delta_{i-1} \cap \Delta_i$ is a weak reducing pair for $1 \leq i \leq n$ and
- (2) $(\bar{V}_1, \bar{W}_1) \subset \Delta_0$ and $(\bar{V}_2, \bar{W}_2) \subset \Delta_n$.

By the assumption that every \mathcal{V} - or \mathcal{W} -face has OEC, each Δ_i has OEC for $0 \leq i \leq n$. Therefore, each Δ_i belongs to a uniquely determined equivalent cluster B^i for $0 \leq i \leq n$ by Lemma 3.31. Since two different equivalent clusters cannot share a weak reducing pair by Lemma 3.29, we conclude $B_{\alpha_1} = B^0 = \dots = B^n = B_{\alpha_2}$, violating the assumption that $B_{\alpha_1} \neq B_{\alpha_2}$.

This completes the proof of Lemma 4.2. \square

Suppose there are two equivalent clusters such that the corresponding equivalent classes determined by Φ are the same. The next lemma gives a sufficient condition that these two equivalent clusters are the same.

Lemma 4.3. *Suppose M is an orientable, irreducible 3-manifold and $(\mathcal{V}, \mathcal{W}; F)$ is a weakly reducible, unstabilized Heegaard splitting of M of genus at least three. Let B_1 and B_2 be equivalent clusters such that $\Phi(B_1) = \Phi(B_2)$ and (\bar{V}_i, \bar{W}_i) the center of B_i for $i = 1, 2$. Suppose if \bar{V}_i (resp. \bar{W}_i) is separating, then the closure of the component of $\mathcal{V} - \bar{V}_i$ (resp. $\mathcal{W} - \bar{W}_i$) missing $\partial \bar{W}_i$ (resp. $\partial \bar{V}_i$) is $S \times I$, where S is a component of $\partial_- \mathcal{V}$ (resp. $\partial_- \mathcal{W}$) for $i = 1, 2$. Then $B_1 = B_2$.*

Proof. Let $\mathbf{H}_i = (\bar{F}_{\bar{V}_i}, \bar{F}_{\bar{V}_i\bar{W}_i}, \bar{F}_{\bar{W}_i})$ be the GHS obtained by weak reduction from $(\mathcal{V}, \mathcal{W}; F)$ along (\bar{V}_i, \bar{W}_i) for $i = 1, 2$. Since \mathbf{H}_2 is equivalent to \mathbf{H}_1 by the assumption that $\Phi(\mathcal{B}_1) = \Phi(\mathcal{B}_2)$, we get the following:

- (1) \bar{F}_{V_2} and \bar{F}_{W_2} are isotopic to \bar{F}_{V_1} and \bar{F}_{W_1} in \mathcal{V} and \mathcal{W} , respectively, and
- (2) \mathcal{B}_1 and \mathcal{B}_2 are of the same type by Lemma 3.15.

By (2), \bar{V}_1 is nonseparating in \mathcal{V} if and only if \bar{V}_2 is nonseparating in \mathcal{V} and the symmetric argument also holds for \bar{W}_1 and \bar{W}_2 in \mathcal{W} .

Suppose \bar{V}_1 is nonseparating in \mathcal{V} , i.e. both \bar{V}_1 and \bar{V}_2 are nonseparating in \mathcal{V} by the previous argument. Since $\bar{F}_{\bar{V}_2}$ is isotopic to $\bar{F}_{\bar{V}_1}$ in \mathcal{V} by (1), considering Corollary 3.2, \bar{V}_2 is isotopic to \bar{V}_1 in \mathcal{V} by (2.11) of Lemma 2.11.

Suppose \bar{V}_1 is separating in \mathcal{V} , i.e. both \bar{V}_1 and \bar{V}_2 are separating in \mathcal{V} . By the assumption, \bar{V}_1 cuts off $S \times I$ missing $\partial\bar{W}_1$ from \mathcal{V} for a component $S \subset \partial_-\mathcal{V}$ of genus $k \geq 1$. Here, the genus of $\bar{F}_{\bar{V}_1}$ is $n - k$ by Corollary 3.2. Let $\tilde{\mathcal{V}}$ be the closure of the component of $\mathcal{V} - \bar{F}_{\bar{V}_1}$ intersecting F . Then $\tilde{\mathcal{V}}$ is a compression body of genus n with negative boundary consisting of the genus k component S and the genus $n - k$ component $\bar{F}_{\bar{V}_1}$, where $\partial_+\tilde{\mathcal{V}} = \partial_+\mathcal{V}$, by Lemma 2.10. By (1), we can isotope $\bar{F}_{\bar{V}_2}$ into $\bar{F}_{\bar{V}_1}$ by an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that $h_1(\bar{F}_{\bar{V}_2}) = \bar{F}_{\bar{V}_1}$. Here, \bar{V}_1 and $h_1(\bar{V}_2)$ are compressing disks in $\tilde{\mathcal{V}}$, i.e. $h_1(\bar{V}_2)$ is isotopic to \bar{V}_1 in $\tilde{\mathcal{V}}$ by Lemma 2.9. Therefore, $h_1(\bar{V}_2)$ is isotopic to \bar{V}_1 in \mathcal{V} by an isotopy g_t defined on \mathcal{V} by Corollary 2.2. Hence, the sequence of isotopies consisting of h_t and g_t gives an isotopy defined on \mathcal{V} that takes \bar{V}_2 into \bar{V}_1 .

Therefore, \bar{V}_2 is isotopic to \bar{V}_1 in \mathcal{V} in any case. Likewise, we conclude \bar{W}_2 is isotopic to \bar{W}_1 in \mathcal{W} . Hence, $(\bar{V}_1, \bar{W}_1) = (\bar{V}_2, \bar{W}_2)$ in $\mathcal{D}(F)$. Therefore, we get $\mathcal{B}_1 = \mathcal{B}_2$ by Lemma 3.29.

This completes the proof. \square

Finally, we reach Theorem 4.4.

Theorem 4.4. *Let $(\mathcal{V}, \mathcal{W}; F)$ be a weakly reducible, unstabilized Heegaard splitting of genus three in an orientable, irreducible 3-manifold M . Then the domain of Φ is the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$, Φ is bijective, and there is a canonically induced function Ω from the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ to the set of the isotopy classes of the generalized Heegaard splittings obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$. The number of components of the preimage of an isotopy class of Ω is the number of ways to embed the thick level contained in \mathcal{V} into \mathcal{V} (or in \mathcal{W} into \mathcal{W}). This means if we consider a generalized Heegaard splitting \mathbf{H} obtained by weak reduction from $(\mathcal{V}, \mathcal{W}; F)$, then the way to embed the thick level of \mathbf{H} contained in \mathcal{V} into \mathcal{V} determines the way to embed the thick level of \mathbf{H} contained in \mathcal{W} into \mathcal{W} up to isotopy and vice versa.*

Proof. By Corollary 3.6, every \mathcal{V} - or \mathcal{W} -face has OEC. Therefore, Lemma 4.2 means every component of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ is an equivalent cluster.

Let \mathcal{B}_1 and \mathcal{B}_2 be equivalent clusters such that $\Phi(\mathcal{B}_1) = \Phi(\mathcal{B}_2)$ and (\bar{V}_i, \bar{W}_i) the center of \mathcal{B}_i for $i = 1, 2$. Considering Lemma 3.27, if \bar{V}_i is separating, then it cannot cut off a solid torus from \mathcal{V} . Moreover, since the genus of F is three, the closure of the component of $\mathcal{V} - \bar{V}_i$ containing $\partial\bar{W}_i$ is a genus two compression body by Lemma 2.3 and Lemma 1.3 of [12]. This means the closure of the other component must be $(\text{torus}) \times I$ if \bar{V}_i is separating. The symmetric argument also holds for \bar{W}_i . Therefore, Lemma 4.3 induces $\mathcal{B}_1 = \mathcal{B}_2$, i.e. Φ is an injective function from the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ to \mathcal{GHS}_F^* / \sim . Considering

Definition 3.30, we conclude Φ is bijective.

Recall that \mathcal{GHS}_F is the set of isotopy classes of the GHSs obtained by weak reductions from $(\mathcal{V}, \mathcal{W}; F)$. Since all GHSs in an equivalent class of \mathcal{GHS}_F^* / \sim correspond to the same isotopy class, there is a function Ω from the set of components of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ to \mathcal{GHS}_F canonically induced from Φ .

Let us consider the preimage $\Omega^{-1}([\mathbf{H}])$ for an isotopy class $[\mathbf{H}] \in \mathcal{GHS}_F$. Let \mathcal{B}_1 and \mathcal{B}_2 be two equivalent clusters in $\Omega^{-1}([\mathbf{H}])$, (V_i, W_i) a weak reducing pair in \mathcal{B}_i for $i = 1, 2$, and \mathbf{H}_i the GHS obtained by weak reduction along (V_i, W_i) for $i = 1, 2$.

If $\bar{F}_{V_2} := \text{Thick}(\mathbf{H}_2) \cap \mathcal{V}$ is not isotopic to $\bar{F}_{V_1} := \text{Thick}(\mathbf{H}_1) \cap \mathcal{V}$ in \mathcal{V} , then $\Phi(\mathcal{B}_1) = [\mathbf{H}_1]^* \neq [\mathbf{H}_2]^* = \Phi(\mathcal{B}_2)$ and therefore $\mathcal{B}_1 \neq \mathcal{B}_2$.

Suppose \bar{F}_{V_2} is isotopic to \bar{F}_{V_1} in \mathcal{V} . Without changing the equivalent classes $[\mathbf{H}_1]^*$ and $[\mathbf{H}_2]^*$, we can assume (V_i, W_i) is the center of \mathcal{B}_i for $i = 1, 2$ by Lemma 3.25.

We claim that $\mathcal{B}_1 = \mathcal{B}_2$. For the sake of contradiction, suppose \mathcal{B}_1 and \mathcal{B}_2 are different. By the assumption, we can isotope \bar{F}_{V_2} into \bar{F}_{V_1} by an isotopy $h_t : \mathcal{V} \rightarrow \mathcal{V}$, $t \in [0, 1]$ such that $h_1(\bar{F}_{V_2}) = \bar{F}_{V_1}$. Let \mathcal{V}' be the closure of the component of $\mathcal{V} - \bar{F}_{V_1}$ intersecting F . Then \mathcal{V}' is a genus three compression body with at least one negative boundary component \bar{F}_{V_1} whose genus is two by Corollary 3.2 and Lemma 2.10 and V_1 and $h_1(V_2)$ are compressing disks in \mathcal{V}' , where $\partial_+ \mathcal{V} = \partial_+ \mathcal{V}'$.

If $\partial_- \mathcal{V}'$ is connected, then every separating compressing disk in \mathcal{V}' cuts off a solid torus from \mathcal{V}' and also does in \mathcal{V} . Therefore, considering that each V_i belongs to the center of \mathcal{B}_i for $i = 1, 2$, Lemma 3.27 forces V_1 and V_2 to be nonseparating disks in \mathcal{V} , i.e. ∂V_1 and ∂V_2 are nonseparating in $\partial_+ \mathcal{V} = \partial_+ \mathcal{V}'$. This means V_1 and $h_1(V_2)$ are nonseparating compressing disks in \mathcal{V}' . But there is a unique nonseparating disk in \mathcal{V}' up to isotopy by Lemma 2.8, i.e. $h_1(V_2)$ is isotopic to V_1 in \mathcal{V}' . Therefore, $h_1(V_2)$ is isotopic to V_1 in \mathcal{V} by an isotopy g_t defined on \mathcal{V} by Corollary 2.2. Hence, the sequence of isotopies consisting of h_t and g_t gives an isotopy defined on \mathcal{V} taking V_2 into V_1 . But this means the centers of \mathcal{B}_1 and \mathcal{B}_2 intersect each other, violating the assumption that each component of $\mathcal{D}_{\mathcal{V}\mathcal{W}}(F)$ is an equivalent cluster itself. If $\partial_- \mathcal{V}'$ is disconnected, then V_1 and $h_1(V_2)$ are isotopic in \mathcal{V}' by Lemma 2.9. Hence, we also get a contradiction similarly.

Therefore, $\mathcal{B}_1 = \mathcal{B}_2$, i.e. $\Phi(\mathcal{B}_1) = [\mathbf{H}_1]^* = [\mathbf{H}_2]^* = \Phi(\mathcal{B}_2)$ and therefore $\text{Thick}(\mathbf{H}_2) \cap \mathcal{W}$ is also isotopic to $\text{Thick}(\mathbf{H}_1) \cap \mathcal{W}$ in \mathcal{W} .

Hence, we only need to count the number of ways to embed the thick level contained in \mathcal{V} into \mathcal{V} in order to count the number of elements of $\Omega^{-1}([\mathbf{H}])$.

This completes the proof. \square

ACKNOWLEDGEMENTS. The author gives thanks to the referee for a number of helpful suggestions for improvement in the article. The author is grateful to Professor Jongil Park and the other professors of the SNU Mathematical Sciences Division for giving me the opportunity to be a postdoctoral researcher for four years. This research was supported by BK21 PLUS SNU Mathematical Sciences Division.

References

- [1] D. Bachman: *Connected sums of unstabilized Heegaard splittings are unstabilized*, Geom. Topol. **12** (2008), 2327–2378.
- [2] A. Ido, Y. Jang and T. Kobayashi: *Heegaard splittings of distance exactly n* , Algebr. Geom. Topol. **14** (2014), 1395–1411.
- [3] J. Kim: *On critical Heegaard splittings of tunnel number two composite knot exteriors*, J. Knot Theory Ramifications **22** (2013), 1350065, 11 pp.
- [4] J. Kim: *On unstabilized genus three critical Heegaard surfaces*, Topology Appl. **165** (2014), 98–109.
- [5] J. Kim: *A topologically minimal, weakly reducible, unstabilized Heegaard splitting of genus three is critical*, Algebr. Geom. Topol. **16** (2016), 1427–1451.
- [6] M. Lustig and Y. Moriah: *Closed incompressible surfaces in complements of wide knots and links*, Topology Appl. **92** (1999), 1–13.
- [7] H. Masur and S. Schleimer: *The geometry of the disk complex*, J. Amer. Math. Soc. **26** (2013), 1–62.
- [8] D. McCullough: *Virtually geometrically finite mapping class groups of 3-manifolds*, J. Differential Geom. **33** (1991), 1–65.
- [9] K. Morimoto: *On Heegaard splittings of knot exteriors with tunnel number degenerations*, Topology Appl. **196** (2015), 719–728.
- [10] R. Gelca: *Theta functions and knots*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2014.
- [11] T. Saito, M. Scharlemann and J. Schultens: *Lecture notes on generalized Heegaard splittings*, arXiv:math/0504167v1.
- [12] M. Scharlemann and A. Thompson: *Heegaard splittings of $(\text{surface}) \times I$ are standard*, Math. Ann. **295** (1993), 549–564.
- [13] M. Scharlemann and A. Thompson: *Thin position for 3-manifolds*, AMS Contemp. Math. **164** (1994), 231–238.

BK21 PLUS SNU Mathematical Sciences Division
 Seoul National University
 1 Gwanak-ro, Gwanak-gu, Seoul 08826
 Republic of Korea

[Current Address:]
 Dong-gu Office
 147, Dong-gu Office Street, Dong-gu
 Daejeon 34691
 Republic of Korea
 e-mail: pibonazi@gmail.com